

INDEX

- Abacoan Stage, defined, 737
 Abacoan Zone NC2, 741
 Acoustic "blanking", 529
 Albian Claystone, 587
 red sediment deposits, Blake Nose, 84
 Albian-Campanian stratigraphic hiatus, 490, 492
 Albian-Turonian transgression, 942
 Algal stromalites, 414
 Aluminum content, red limestone, 505
 variegated and red claystone, 507
 Aluminum concentration in sediments, 576
 Ammonites, 10, 80, 170, 181, 193
 Aulacostarphanus mutabilis Zone, 762
 Pectinatites pectinatus Zone 762, 763
 See Also: Aptychi
 Ammonitico Rosso facies, 465, 957
 similarity of Atlantic Jurassic section to, 10
Amphirhopalum ypsilon Zone, 194, 866
 Analyses of sediments, methods, 478
 Angola Basin, C₁-C₇ hydrocarbons, 651
 Antarctic glaciations, effect on sea level, 202
 Antarctic Bottom Water Current, 153
 Appalachian piedmont, source of clay fraction in
 variegated ooze, 84
 Aptychi, assemblage of, 899
 Early Cretaceous, 899
 Lamellaptychus altus, new species, 904
 angulocostatus planus, new subspecies, 905
 bahamensis, new species, 903
 challengeri, new species, 904
 postbermudensis, new species, 904
 stellariformis, new species, 904
 subseranonis, new species, 904
 transitorius, new species, 903
 Late Jurassic, 899
 Aragonite, occurrence of, 519
 Arctic Bottom Water, effect on circulation patterns,
 103
 Argusian Stage, defined, 737
Arkangelskiella cymbiformis Zone, 93, 96, 98, 710
 Ascidian spicules, 810
 Atlantic Ocean, Jurassic spreading rates, 8
 Atlantic coastal plain, 941
 Atlantic margin, block faulting, 941
 rifting, 941
Aulacoastephanus mutabilis Zone, 762
 Bahama channels, marine sediment source, 181
 source of reworked Cretaceous fossils, 191
 Bahama Banks, 529
 Bahama Outer Ridge, 153
 Barremian stratigraphic hiatus, 953
 limestone, depositional environment, 80
 Basement reflector, 540
Bekoma bidartensis Zone, 93, 99, 865
 Benthic foraminifers, Miocene, 190
 Site 388, 36
 Tertiary, 36
 zonation
 Epistomina mosquensis Zone, 181, 911
 Gaudyrina ex. gr. *dividens* Zone, 664, 669
 Gavelinella barremiana-Lenticulina (M.) sigali
 Zone, 97, 664, 673, 676
 Bermudan Stage, defined, 738
Biorbifera johnewingii Zone, defined, 890
Biorbifera johnewingensis-Deflandrea vestita zones, 890
 Biostratigraphy, Blake Nose, 671
 Jurassic, lithostratigraphy, 941
 Lower Cretaceous, 941
 Site 388, 34, 657
 Site 389, 89, 657
 Site 390, 89, 657
 Site 391, 179, 660
 Site 392, 351, 657
 Tertiary, 943
 Upper Cretaceous, 942
Biscutum dimorphosum Zone (NP 3), 98, 710
 defined, 704
 Bitumens, 645
 hydrocarbon fraction of, 645
 Black sea, hydrocarbon samples from sediments of, 651
 Blake-Bahama Basin, depth of deposition of
 sediments, 951
 Blake Escarpment, 198, 539
 bottom currents along, 108
 erosion, 338
 reef-lagoon facies along, 70, 337
 sediment sampling from, 664
 Blake Nose, 69, 105, 540, 550, 605, 615, 657, 663, 673
 676
 Albian red sediment deposits, 84
 biostratigraphy, 671
 Cretaceous reef building, 356
 Objectives and results, 7
 sedimentary history, 85
 subsidence of, 951
 Blake Plateau, 358, 599, 663, 673
 marine sediment source, 181, 194, 203
 reflectors, 540
 sediment transport across, 85
 source of reworked Cretaceous fossils, 191
 Blake-Bahama Abyssal Plain, 529
 sediment sampling from, 664
 Blake-Bahama Basin, 198, 529, 599, 605, 615, 617,
 639, 660, 663
 age of sediments, 587
 clay mineralogy, 949
 Cretaceous black shales from, 635
 Cretaceous claystones in post-Barremian sediments
 of the, 593
 C₁-C₇ hydrocarbons, 651
 depositional events, 194
 fauna, 951
 geological history of, 625
 "Magnetic Quiet Zone," 201
 Miocene reflectors, 529
 objectives and results, 8
 sediment accumulation rate, 202
 sedimentological features, 949
 Site 391, 153
 Site 393, 395

- Site 394, 395
 turbidites, upper Miocene, 201
 Blake-Bahama Outer Ridge, 550
 Block faulting, Atlantic margin, 941
Bonetocardiella paulaworstellae, new
 Calcisphaerulidae species, 912
Bonetocardiella williamsonii, new,
 Calcisphaerulidae species, 912
 Bottom currents, 530
 along Blake Escarpment, 108
 effect on sediment accumulation, 108
Brianconnais series compared to Leg 44 sediments, 571
Brianconnais Zone, 954
 Budden Canyon formation, 672, 676
 Bulk geochemical data, 579
Buryella clinata Zone, 93, 99, 865
 Calcareous microfossils, Eocene/Paleocene boundary,
 93
 Tertiary/Cretaceous boundary, 93
Calcicalathina oblongata Zone, 714
Calcicalathina oblongata/Lithraphidites bollii Zonal
 interval, 193
Calcisphaerulidae, Bonetocardiella paulaworstellae,
 new Species, 912
Bonetocardiella williamsonii, new species, 912
 systematic description of, 912
 Upper Jurassic, 911
 Calcite overgrowth, nannofossils, 733
Calocyclus costata Zone, 180, 194, 866, 867
 Campanian-Albian angular unconformity, 7
 sediments, stratigraphic hiatus, 106
 stratigraphic hiatus, 707
 unconformity, 541
Campylosphaera eodola Subzone, 782
 Canadian Atlantic margin, 673
 Canadian Atlantic Shelf, 676
 Canaverlain Stage, defined, 737
 Cape Basin, C₁-C₇, hydrocarbons, 651
 Carbohydrate residues in sediments, 585, 653
 Carbon, geochemistry of 605
 Carbon isotopic composition, Cretaceous organic
 matter, 585
 Carbon preference index (CPI), 627
 Carbon-carbonate analyses, 15
 Carbon-13 content, humic acids, 589
 kerogen, 606, 611
 lipid fraction, 606, 611
 Carbonaceous shales, source of, 585
 Carbonate, iron concentration in, 562
 manganese concentration in, 557, 561
 strontium concentration in, 557, 559
 Carbonate compensation depth, effect on microfossil
 assemblages, Site 388, 657
 Carbonate sediments, Site 391, diagenesis of, 478
 Carbonate, Dissolution Depth, 41
 Carbonate, trace elements in, 557
Caryian Zone NC 1, 741
 Caryian Stage, defined, 737
 Cenozoic nannofossils, 716
 radiolarian, 865
 Cenozoic sedimentation, Leg 44, 8
 Chemical analyses of shipboard lubricants, 645
 Chert, replacement type associated with lithified
 Paleocene limestone, 85
 Chert layer, Eocene, 540
Chiasmolithus gigas Subzone, 782, 807
bidens Subzone, 782
danicus Zone (NP 3), 704, 710
Chiastozygus litterarius/Parhabdololithus angustus
 zonal interval, 193
Chiphragmalithus quadratus Zone, 704
 Chlorins, 585
 mass spectrometric analyses of, 640
 Site 391, 639
 Chloroformic extracts, 596
 Chromium, concentration in sediments, 579
 Clathrates, 5
Clavatipollenites Zone, defined, 892
 Clay mineralogy, Blake-Bahama Basin, 949
 Clay minerals, 509
 Coarse fraction of sediments, Site 390, 484
 Site 391, 488
 Cobalt, concentration in sediments, 579
 Column chromatography analyses, 617
 Concentration in sediments, aluminum, 576
 chromium, 579
 cobalt, 579
 copper, 579
 iron, 575
 manganese, 576
 nickel, 579
 titanium, 576
 zinc, 579
 Coniacian regression, 942
 Continental offshore stratigraphic test, 941
 Contour current, 489, 493, 529
Conusphaera mexicana Zone, 716
 defined, 768
 Copper, concentration in sediments, 579
 native, 28
Corbisema toxema, new silicoflagellate species, 815
triacantha convexa, new silicoflagellate species, 815
triacantha zone, 809, 811, 812
Corbisema? xenica, new species, 816
Craspedodiscus cocinodiscus Zone, 809, 811
 Cretaceous, Nannofossil Stratigraphy, 731
 black shales, 8
 Blake-Bahama Basin, 635
 depositional environment, 627
 source rock potential of, 635
 southern and eastern Atlantic area, 587
 lipid distribution determination from, 591
 clays as sapropelitic black shales, 597
 claystones, organic chemistry of, 593
 in post Barremian sediments, 593
 European nannofossil stages, 733
 foraminifers, 352
 organic matter, carbon isotopic composition, 585
 ostracodes, Site 390, 921
 Site 392, 921
 reef-building, Blake Nose, 356
 reef complex, 105, 356, 663
 seismic evidence for, 69, 108, 337
 stratigraphic units, radiometric ages of, 738
 variegated and red claystones, geochemistry, 505
 Cretaceous-Jurassic limestones, Site 391, environment
 of deposition of, 465
 Cretaceous-upper Jurassic nannofossils, 193
 Site 391, 181
 Cretaceous-Jurassic boundary, dinoflagellates, 891
 Cretaceous-Paleocene boundary, stratigraphic hiatus,
 106
 Cretaceous-Tertiary boundary, 559
Cretarhabdus crenulatus Zone, 193, 714
quadratus, new nannofossil combination, 764

- Cruciplacolithus tenuis* Zone (NP 2), 98, 710, 778, 779, 807, 808
defined, 704
- Cryptocapsella tetrapera* Zone, 866
- Cuba, tectonic uplift in middle Miocene, 202
- Cuche Formation, 674, 676
- Cuchon section, 955
- Current, Tethys, 743
- Cutin composition of lower plants, 621
- Cyclagelosphaera bergeri*, new nannofossil species, 746
reinhardtii, new combination, 746
- Cyrtocapsella tetrapera Zone, 180, 194
- C₁-C₇ hydrocarbons, Angola Basin, 641
Blake-Bahama Basin, 651
Cape Basin, 651
- Debris flow, 530
- Deflandrea echinoidea* Zone, 893
vestita Zone, 892, 893
defined, 892
vestita/Retitricolpites georgensis (Psilaticolporites) zones, 894
- Degree of organic diagenesis (DOD), 617
- Depositional bedding, 27
- Depositional environment, Barremian limestone, 80
Cretaceous black shale episode, 627
fenestral limestone, 437
oolitic limestone, 349, 424
Site 390, 99
Site 391, 181
Site 392, 353
diagenesis, 356
skelmoldic limestone, 416
western North Atlantic, 958
- Depositional events, Blake-Bahama Basin, 194
- Depth of burial, effect on early gas generation, 632
oil generation, 632
- Depth of deposition of sediments, Blake-Bahama 951
- Depth/age profile, tetrapyrrole pigments, 639
- Desmosponges, 807
- Diagenesis, carbonate sediments, Site 391, 478
depositional environment, Site 392, 356
humic acid, 589
organic matter in Cretaceous black claystones, 171
- Diagenetic environment, skelmoldic limestone, 419
- Diagenetic history, fenestral limestone, 439
oolitic limestone, 425
skelmoldic limestone, 417
- Dictyochoa deflandrei lobata*, new silicoflagellate species, 785
- Dictyochoa hexaxantha* Zone, 783, 787
spinosa Subzone, 779, 814
spinosa Zone, 783
- Dinoflagellates
Biorbifera johnewingii-Deflandrea vestita zones, 890
Biobifera johnewingii Zone, defined, 890
Cretaceous-Jurassic boundary, 891
- Deflandrea echinoidea* Zone, 893
vestita Zone, 892, 893
defined, 892
vestita/Retitricolpites georgensis (Psilaticolporites) zones, 894
- Druggidium apicopaucina* Zone, defined, 891
deflandrei Zone, defined, 891
rhabdoreticulatum Zone, 890
defined, 891
- Odontochitina operculata* Zone, 890, 891
defined, 891
- Oligosphaeridium complex* Zone, 890, 891
- Phobericysta neocomica* Subzone, 890, 891
source of carbohydrates in sediments, 653
- subtilisphaera perlucida* Subzone, 890, 892
- Trithyrodinium suspectus/complexipollis* zones, 893
- Discoaster berggrenii* Subzone, 810
calcaris Zone (NN10), 38, 707
druggi Zone (NN2), 712
exilis Zone, 814
hamatus Zone (NN9), 38, 41, 707
kugleri subzone, 810
lodoensis Zone (NP13), 98, 709, 710, 775, 782, 783, 808, 814
mohleri Zone (NP7), 98, 710, 776, 783
multiradiatus Zone (NP 18), 710, 776
neohamatus Zone, 807
nobilis Zone, 776, 778, 783
quinqeramus Zone (NN 11), 38, 41, 180, 707, 711
strictus Subzone, 782
sublodoensis Zone (NP 14), 98, 704, 709
tani nodifer Zone (NN 16), 703, 708
tani s.l.-*Sphenolithus radians* Zone, 704
t. Chiasmolithus ratio, 775, 781
- Discoasteroides kuepperi* Subzone, 775
- Distephanus crux loeblichii*, new silicoflagellate sub-species, 817
? acanthicus, new silicoflagellate species, 816
triotus, new silicoflagellate species, 818
- Diversity of nannofossils, 745
- Dorcadospyris alata* Zone, 180, 194, 866
- Druggidium apicopaucium* Zone, defined, 891
deflandrei Zone, defined, 891
deflandrei/Ephedripites multicostatas Zones, 894
rhabdoreticulatum Zone, 890
defined, 891
- Duck Creek Formation, 673
- Early gas generation, depth of burial effect on, 632
- Early Cenozoic foraminifers, 95
- Early Cretaceous aptychi, 899
nannofossils, evolution of, 742
western and eastern Atlantic, correlation of, 899
- Effective carbon in sediments, defined, 599
- Emiliania annula* Subzone, 775
huxleyi Zone, 778
- Environments of deposition, Cretaceous-Jurassic limestone, Site 391, 465
- Eocene, chert layer, 540
nannofossils, 98, 708
- Eocene-Paleocene boundary, calcareous microfossils, 93
- Ephedripites multicostatas* Zone, defined, 892
Retitricolpites georgensis Zone, 890
- Epistomina mosquensis* Zone, 181, 911
- Erosion, Blake Escarpment, 338
- Erosion hypothesis, lower continental rise hills, 43
- Etching, nannofossils, 739
- European nannofossil stages, Cretaceous, 733
- Evolution, nannofossils, early Cretaceous, 742
- Fauna, Blake-Bahama Basin, 951
- Feldspar, 510
- Fenestral limestone, 432, 489, 951
depositional environment, 437
diagenetic history, 439

- lithology, 432
- sedimentary structures, 433
- Fertility, nannofossil, 742
- Foraminifers, Cretaceous, 352
 - early Cenozoic, 95
 - Jurassic red limestone, 193
 - Lower Cretaceous, 97
 - Miocene, 189, 192
 - Pleistocene, 36, 192
 - Quaternary, 186
 - Site 388, 35
 - Sites 389 and 390, 93
 - Site 391, 181
 - Site 392, 351
 - upper Cenozoic, 95
 - Upper Cretaceous, 96, 352
 - Zonation, 11
 - Globigerinelloides blowi* Zone (LC8), 352
 - Globigerina eugubina* (P1)-*Globorotalia pseudomenardii* (P. 4) zones, 93
 - Globigerina eugubina* Zone, 96
 - Globigerinelloides algerianus* Zone (LC 11), 93, 97, 353, 664, 668, 670, 674, 675
 - /Ticinella breggiensis* zones, 676
 - blowi* Zone, 93, 97, 657, 664, 667, 676
 - /Ticinella breggiensis* zones, 674
 - Globorotalia calida* Zone (N 23), 180
 - fohsi* Zone (N 12), 180
 - formosa* (P. 7)-*Globorotalia lehneri* (P. 12) zones, 93
 - kugleri* Zone (N 4), 181, 190
 - periperoacuta* Zone (N 10), 180
 - pleisotumida* Zone (N 17), 180, 189
 - sicanus* Zone (N 9), 180
 - truncatulinoidea* (N 22), 36, 180
 - Globotruncana arca* zone, 97
 - calcarata* Zone (UC 12), 93, 97, 352, 671
 - contusa* Zone (UC 16), 93
 - gansseri* Zone, 96
 - mayaroensis* Zone, 659
 - scutilla* Zone (UC 13), 93, 97
 - tricarinata* Zone, 97
 - Hedbergella sigali* Zone (LC8), 181, 664, 666
 - Schackoia cabri* Zone, 352
 - cabri-Globigerinelloides algerianus* Zone, (LC 10-11), 353
 - Sphaeroidinellopsis subdehiscens* Zone (N 13), 180
 - Ticinella breggiensis* Zone (LC17), 352, 664, 670
 - primula* Zone (LC 15-16), 93, 97, 353, 664, 669, 671
 - Ticinella primula-Ticinella breggiensis* Zone, 673, 676
- Gas chromatography, 587, 625
 - saturated fraction of hydrocarbons, 596
- Gas clathrates, Site 388, 33
- Gaudryina* ex. gr. *dividens* Zone, 664, 669
- Gavelinella barremiana-Lenticulina sigali zones, 97, 664, 673, 676
- Geochemical analyses, organic carbon, 585
- Geochemistry, carbon, 605
 - Cretaceous variegated and red claystones, 505
 - Site 388, 33
 - Site 390, 86
 - Site 391, 171
 - Site 392, 349
- tetrapyrrole pigments, 639
- Upper Jurassic red limestone, 504
- Geologic cross-section across the Blake Nose, 108
- Geological history, Blake-Bahama Basin, 625
- Geological setting, north Tethyan margin, 953
- Gephyrocapsa caribbeanica* subzone, 807
- Gephyrocapsa oceanica-Emiliana huxleyi* zonal interval, 38, 41, 180, 353, 703, 707, 711
- Globigerinelloides blowi* Zone (LC9), 352
- Globigerina eugubina* P. 1)-*Globorotalia pseudomenardii* (P. 4) Zone, 93
- Globigerina eugubina* P. 1) Zone, 93
- Globigerinelloides algerianus* Zone (LC11), 93, 97, 353, 664, 668, 670, 674, 675
- Globigerinelloides algerianus-Ticinella breggiensis* Zone, 676
- Globigerinelloides blowi* Zone (LC9), 93, 97, 657, 664, 667, 676
- Ticinella breggiensis* zones, 674
- Globorotalia calida* Zone (N23), 180
- fohsi* Zone (N12), 180
- formosa* (P. 7)-*Globorotalia lehneri* (P. 12) Zones, 93
- kugleri* Zone (N 4), 181, 190
- periperoacuta* Zone (N 10), 180
- peripheroronda* Zone (N 9), 180
- pleisotumida* Zone (N 17), 180, 189
- sicanus* Zone (N 9), 180
- truncatulinoidea* (N 22), 36, 180
- Globotruncana arca* Zone, 97
- calcarata* Zone (UC 12), 93, 97, 352, 671
- contusa* Zone (UC16), 93
- gansseri* Zone, 96
- mayaroensis* Zone, 659
- scutilla* Zone (UC 13), 93, 97
- tricarinata* Zone (UC 13), 97
- Goethite, 44
- Gorringe Bank, 675, 676
- Graham Interval, 154
- Grand Banks, 674
 - Shelf area, Jurassic biozonation, 191
- Gravity transport, Site 391, 162, 201
- Grayson formation, 673
- Great Abaco Fracture Zone, 203
- Great Abaco Canyon, 203, 205, 358
- Great Bahama Banks, pelletal sand environment, 346
- Green River shale compared to Cretaceous black shales, 636
- Gulf of Mexico, slumped beds, 27
- Hamius circumradiatus*, new nannofossil combination, 746
- Hatteras Abyssal Plain, 676
 - sediment sampling from, 664
- Heavy minerals, 510
- Hedbergella sigali* Zone (LC 8), 181, 664, 666
- Helicosphaera ampliapertura* Zone (NN4), 181, 712, 809, 813
- ampliapertura/Sphenolithus heteromorphus* zones (NN4/5, 713
- reticulata* Zone, 781
- Hexactinellids, 807
- Hole 386, sediment accumulation rates, 504
- Hole 387, sediment accumulation rates, 504
- Hole 390, isotopic data, 567
- Hole 390A, isotopic data, 567

- Hole 391A, radiolarians, 865
Hole 391C, *Aptychus* assemblages, 899
isotopic data, 568
paleomagnetic results, 524
sediments, 523
Hole 392A, isotope data, 567
See also: Sites 386 - 392
Horizon A, 8, 10, 530, 537, 550
Horizon Beta, 8, 10, 537
Howlandian Stage, defined, 738
Hudson submarine canyon, marine sediment source, 205
Humic acids, 585, 591, 593, 597
carbon-13 content, 589
diagenesis, 589
Site 391, sources of, 587
Hydrocarbons, analyses of, 617
bitumen fraction, 645
Black sea sediments, 651
concentration, 585
gas, potential source for, 585
generation, 618
mass-spectrometric analyses, 645
maturation, 594
Hydrogen/carbon ratios, 627
Incertae sedies, *Rocella schraderi*, new species, 788
Rocella gelida, new combination, 788
Index species, nannofossils, criteria for, 733
Interstitial gas, 6
Interstitial water, Site 388, 34
Site 390, 86
Site 391, 171
Iron and manganese minerals, 515
Iron, concentration in carbonate, 562
concentration in red limestone, 505
concentration in sediments, 575
Ismolithus recurvus Zone, 781
Isotope balance, 568
Isotopic data, Hole 390, 567
Hole 390A, 567
Hole 391C, 568
Hole 392A, 567
Jurassic biozonation, Grand Banks Shelf area, 191
Scotia Shelf, 192
lithostratigraphy, 941
red limestone, 193
spreading rates, Atlantic, 8
western and eastern Atlantic, correlation, 902
Keathley magnetic anomaly sequence, 27, 154
Kelvin Seamounts, magnetic profiles, 24
Kerogen, 585, 587, 593, 597, 617, 622, 627, 645
carbon-13 content, 606, 611
in lithified sediments, 589
maceral composition, 602
maturation profile, 629
La Plisse Section, 956
Lamellaptychus altus, new aptychi species, 904
angulocostatus planus, new aptychi subspecies, 905
bahamensis, new aptychi species, 903
challengeri, new aptychi species, 904
postbermudensis, new aptychi species, 904
stellariformis, new aptychi species, 904
subseranonis, new aptychi species, 904
transitorius, new aptychi species, 903
Lamination, 345
Lamprocyrtis haysi Zone, 180
Late Cretaceous, variegated sediments, origin of, 515
Late Jurassic, aptychi, 899
Level of organic metamorphism (LOM), 599, 600
Light hydrocarbons, 651
Limestone, aluminum content, 505
fenestral, 489, 951
geochemistry, red limestone, 504
iron content, 505
manganese content, 505
red stained, 440
sedimentation rate, 503
Site 391, environment of deposition, 465
skelmoldic, 489
titanium content, 505
Limonite pisolites, 356
Lipids, 585, 587, 591, 618
distribution, Cretaceous black shales, 591
carbon-13 content, 606, 611
Site 391, sources of, 587
Lithology, fenestral limestone, 432
oolitic limestone, 421
Site 388, 30
Sites 389 and 390, 74
Site 391, 160
Site 392, 339
Site 393, 397
Site 394, 397
skelmoldic limestone, 414
Lithostratigraphy, Jurassic, 941
Lower Cretaceous, 941
Tertiary, 943
Upper Cretaceous, 942
Lithraphidites bolli Zone, 714
praequadrites, new nannofossil species, 749
quadratus Zone, 93, 98, 710, 741
Lower continental rise hills, 605, 615, 657
construction hypothesis of, 43
erosion hypothesis of, 43
objectives and results of drilling, 5
origin of, 27, 43
Site 388, 23
slump deformation hypothesis, 44
Lower Cretaceous, foraminifers, 97
lithostratigraphy and biostratigraphy, 941
nannofossils, 98, 710, 762
stratigraphic hiatus, 98
Lower Paleocene, nannofossil zonation, 704
Lynchnocanoma elongata Zone, 181, 194, 867
Maceral composition, kerogen, 602
Maestrichtian stratigraphic hiatus, 96
-Paleocene stratigraphic hiatus, 492
Magellanian Stage, defined, 737
Magnesium content, variegated and red claystone, 508
Magnetic profiles, Kelvin Seamounts, 24
Scotian Shelf, 24
“Magnetic quiet zone”, 5, 8, 24, 153, 395, 663
Magnetostratigraphic events, 735
Manganese, concentration in carbonate, 557, 561
concentration in red limestone, 505
concentration in sediments, 576
concentration in variegated and red claystone, 506
nodules 7, 559, 807
relationship of nodules to sedimentation rate, 75
Sites 389 and 390, 74
Manganiferous siderite, 581
Maridale formation, 673
Marine bottom topography, Western North Atlantic, 958

- Marine sediments, Bahama channels, 181
 Blake Plateau, 181, 194, 203
 Hudson Submarine Canyon, 205
 Tongue of the Ocean, 181
- Markalius inversus* Zone (NP1), 93, 98, 710
 defined, 706
- Markalius inversus* (NP1)-*Discoaster mohleri* (NP7)
 zones, 93
- Marly limestone, Site 391, 463
- Marthasterites tribrachiatus* Zone (NP 12), 93, 98, 710
Discoaster lodoensis zonal Interval (NP 12/13, 710)
- Mass spectrometry analyses, 587, 610
 chlorins, 640
 hydrocarbons, 645
 nickel porphyrins, 641
 silylated hydroxyaluminum derivatives, 639
 tetrapyrrole pigments, 639
- Maturation, hydrocarbons, 594
- Merrittian Stage, defined, 737
- Mesocena diodon nodosa*, new silicoflagellate
 subspecies, 818
 ? *connudata*, new species, 786
ovata, new species, 786
- Mesozoic nannofossils, 731
- Metalliferous deposits, 5, 28
- Methods of X-ray mineralogy, 515
- Micrantholithus hoschulzi* Zone, 93, 98, 353, 657, 711
Chiatozygus litterarius zonal interval, 193
- Microfacies, Type I defined, 480
 Type II defined, 480
 Type III defined, 480
 Type IV defined, 481
 Type V defined, 481
 Type VI defined, 481
- Microfaults, 465
- Micrograin size analyses, Site 388, 482
 Site 390, 485
 Site 391, 488
 Site 392, 490
- Microkarst formation, Site 392, 349
- Micronodules, 949
- Micula mura* Zone, 93, 96, 98, 710
- Middle Eocene nannofossil zonation, 703
- Mineral carbon, 593
- Mineralogy, Leg 44, sediments, 509
 Site 388, 482
 Site 390, 485
 Site 391, 488
 Site 392, 489
- Miocene, benthic foraminifers, 190
 carbonate turbidites, 8
 foraminifers, 189, 192
 Nannofossils, 707, 711
 reflectors, 529
 turbidites, 530
- Miocene/Cretaceous sediments, 643
 stratigraphic analyses, Site 391, 625
 stratigraphic hiatus, 203, 630
 unconformity, 10
- Moler Formation, 787
- Monosaccharides, paper chromatography analyses of,
 653
- Mud dunes, formation of by contour currents, 5
n-alkanes, 596, 605, 611, 618
 sources of, 587
n-fatty acids, 588
- Nannofossils, calcite overgrowth on, 733
 Cenozoic, 716
 Cretaceous biostratigraphy, 731
 Cretaceous-Upper Jurassic, 733
Cretarhabdus quadratus, new combination, 764
 Criteria for index species, 733
Cycligelosphaera reinhardtii, new combination, 746
Discoaster/Chiasmolithus ratio, 775, 781
 Diversity of, 745
 Early Cretaceous, 742
 Eocene, 98, 708
 etching, 739
 Lower Cretaceous, 98, 710
 Mesozoic, 731
 middle Eocene, 703
 Miocene, 707, 711
 overgrowth, 733, 739
 Paleocene, 98, 710
 preservation of, 739
 Quaternary, 707, 711
 Site 382, 775
 Site 384, 775
 Site 385, 778
 Site 386, 780
 Site 387, 782
 Site 388, 37, 707, 807
 Site 389, 707, 807
 Sites 389 and 390, 97
 Site 390, 707, 739, 807
 Site 391, 711, 741, 768, 808
 Site 392, 353, 707, 810
 Site 393, 711
 Site 394, 711, 810
 systematic descriptions, 746
 Tertiary-Cretaceous boundary, 98, 710
 Upper Cretaceous, 98, 710
 Upper Jurassic-Lower Cretaceous, 762
 zonation, 11, 703
Abacoan Zone NC 2, 741
Arkangelskiella cymbiformis, 93, 96, 98, 710
Biscutum dimorphosum (NP3), 98, 704, 710
Calicalathina oblongata Zone, 714
 - *Lithraphidites bollii* zonal interval, 193
Campylosphaera eodela Subzone, 782
Caryian Zone NC1
Chiasmolithus danicus (NP 3), 704, 710
bidens Subzone, 782
gigas, 782, 807
Chiatozygus litterarius-Parhabdolithus angustus
 zonal interval, 193
Chiphragmalithus quadratus Zone, 704
Conusphaera mexicana Zone, 716
 defined, 768
Cretarhabdus crenulatus Zone, 193, 714
Cruciplacolithus tenuis Zone (NP2), 98, 710,
 778, 779, 807, 808
 defined, 704
Discoaster berggrenii Subzone, 810
Discoaster calcaris Zone (NN10), 38, 707
druggi Zone (NN2), 712
exilis Zone, 814
hamatus Zone, 38, 41, 707
kugleri Subzone, 810
lodoensis Zone, 98, 775, 782, 783, 808, 814
mohleri Zone, (NP 7), 98, 710, 776, 783
multiradiatus Zone (NP 18), 710, 776

- neohamatus* Zone, 807
nobilis Zone, 776, 778, 783
quinqueramus Zone (NN11), 38, 41, 180, 711
strictus Subzone, 782
sublodoensis Zone (NP14), 98, 704, 709
tani nodifer Zone (NN16), 703, 708
tani s.l.-*Sphenolithus radians* Zone, 704
Discoasteroides kuepperi Subzone, 775
Emiliania annula Subzone, 775
huxleyi Zone, 778
Gephyrocapsa caribbeanica Subzone, 807
oceanica-Emiliania huxleyi zonal interval (NN20/21), 38, 41, 180, 353, 707, 711
Helicosphaera ampliapertura Zone (NN4), 181, 712, 809, 813
reticulata Zone, 781
Sphenolithus heteromorphus zones (NN4/5), 713
Ismolithus recurvus Zone, 781
Lithraphidites bollii Zone, 714
quadratus Zone, 93, 98, 710, 714
Markalius inversus Zone (NP1), 93, 98, 710 defined, 706
Markalius inversus-Discoaster mohleri Zones, 93
Marthasterites tribrachiatus Zone (NP12), 93, 98, 710
-Discoaster lodoensis zonal interval (NP12/13), 710
Micrantholithus hoschulzi Zone, 93, 98, 353, 657, 711
Chiatozygus litterarius zonal interval, 193
Micula mura Zone, 93, 96, 98, 710
Nannoconus colomii Zone, 193, 715, 766 defined, 768
Nannotetrina alata Zone (NP15), 98, 703, 709 defined, 703
quadrata Zone, 775, 782, 783
Parahabdolithus angustus Zone, 93, 98, 193, 353, 669, 710, 711, 713
embergeri Zone, 193, 715, 911 defined, 768
Prediscosphaera cretacea Zone, 353, 711
Pseudoemiliania lacunosa Zone (NN19), 38, 41, 180, 707, 711
Reticulofenestra umbilica Zone, 782, 807
-Sphenolithus furcatolithoides zones, 703
Rhabdosphaera inflata Subzone, 782
Sphenolithus belemnoides Zone (NN3), 712, 810
ciperoensis Zone, 781, 788, 810
furcatolithoides Zone (NP16), 93, 98, 703, 708, 709
heteromorphus Zone (NN5), 180, 712, 713, 809, 810, 814
Tetralithus trifidus Zone, 98, 353, 710, 810
Tribrachiatus orthostylus Zone, 808, 814
Triquetrorhabdulus carinatus Zone (NN1), 181, 712
Tubodiscus jurapelagicus Zone, 714
Watznaueria communis Zone, defined, 761
Nannotetrina alata Zone, 98, 703, 709 defined, 703
quadrata Zone, 775, 782, 783
 Native copper, 28
 Naturalistian Stage, defined, 738
Naviculopsis biapiculata, 779, 783
constricta Zone, 776, 783, 786
eobiapiculata, new silicoflagellate species, 787
foliacea tumida, new silicoflagellate subspecies, 820
foliacea Zone, 783, 787, 812, 814
obtusarca, new silicoflagellate species, 865
robusta Subzone, 807, 814
 New England Seamounts, 503
 New species, Aptychi, *Lammelaptychus altus*, 904
bahamensis, 903
challengeri, 904
postbermudensis, 904
stellariformis, 904
subseranonis, 904
transitorius, 903
 Calcisphaerulidae, *Bonetocardiella paulaworstellae*, 912
williamsonii, 912
 Incertae sedis, *Rocella schraderi*, 788
 Nannofossils, *Cyclagelosphaera bergeri*, 746
Lithraphidites praequadrates, 749
 Silicoflagellates, *Corbisema toxseuma*, 815
Corbisema? xenica, 816
Distepahnus triotus, 818
Distephanus? acanthicus, 816
Mesocena ovata, 786
Mesocena? connudata, 786
Naviculopsis eobiapiculata, 787
obtusarca, 865
 New Subspecies
 Aptychi, *Lammelaptychus angulocostatus planus*, 905
 Silicoflagellates, *Corbisema triacantha convexa*, 815
Dictyocha deflandrei lobata, 785
Distephanus crux loeblichii, 817
Mesocena diodon nodosa, 818
Naviculopsis foliacea tumida, 820
 New time scale, paleomagnetic stratigraphy, 155
 Nickel, 28
 concentration in sediments, 579
 porphyrins, 641
 mass spectrometric analyses of, 641
 Norfolk to Site 388, Underway data, 541
 North Tethyan margins, comparison to western North Atlantic, 957
 geological setting of, 953
Odontochitina operculata Zone, 890, 891 defined, 891
 OEP values, 606, 611, 615
 Oil generation, depth of burial effect on, 632
Oligosphaeridium complex Zone, 890, 891
 Ooids, 342, 346, 463
 Oolites, depositional environment of, 349
 Site 392, 343
 Oolitic bed, Schooner Cays, 428
 Oolitic limestone, 421
 depositional environment of, 424
 diagenetic history of, 425
 lithology, 421
 sedimentary structures in, 421
 Organic carbon, 594, 626, 635
 geochemical analyses, 585
 sediments, Sites 391 and 392, 600 defined, 599
 Organic extracts, thermal immaturity evidence
 Organic geochemical analyses, 645, 646
 Cretaceous claystones, 593
 Site 388, 33, 605, 645

- Site 390, 606
 Site 391, 606, 617, 625, 646
 Organic matter, 645
 black claystone, 171, 625
 diagenesis, 171
 low-temperature diagenesis of, 615
 Site 391, petroleum-bearing potential, 617
 Sites 391 and 392, 600
 Organic phosphate, 511
 Ostracodes, Site 390, 921
 Site 392, 921
 Overgrowth on nannofossils, 739
 Paleocene nannofossils, 98, 710
 Paleocene/Eocene boundary, stratigraphic hiatus, 106
 Paleomagnetic results, Hole 391C, 524
 stratigraphy, new time scale for, 155
 Paleotemperature equation, 568
 Palynology, stratigraphic, 888
 Parahabdolithus angustus Zone, 93, 98, 193, 353, 669,
 710, 711, 713
 embergeri Zone, 193, 715, 911
 defined, 768
Pectinatites pectinatus Zone, 762, 763
 Pelletoidal sand environment, Great Bahama Banks, 346
 Petroleum genesis, conditions favoring, 615
 Petroleum-bearing potential, black claystone, 171, 617, 625
 Blake Plateau region, 599
 studies of offshore sources, 585
Phoberocysta neocomica Subzone, 890, 891
Phormocystis striata Zone, 93
 striata striata Zone, 99
 Phosphorite clasts, 74
 Physical properties, Site 388, 34
 Site 390, 86
 Site 391, 171
 Site 392, 349
 Piémont Zone, 957
 Pigment fractions, UV-invisible spectral data of, 639
 Pigments, 585
 Pisolites, 7
 limonite, 356
 Pleistocene, foraminifers, 36, 192
 turbidite deposition, 33
 Polarity sequence, 526
 Potassium content, variegated and red claystones, 508
 Potomac group, 941
Praediscosphaera cretacea Zone, 353, 711
 Preservation, nannofossils, 739
Pseudoemiliana lacunosa Zone (NN 19), 38, 41, 180,
 707, 711
Psilatricolporites Subzone, 890, 892
 Pyrolysis assay method, 594
 FID, 599
 fluorescence, 599
 oil yield, 635
 ratio, 617
 Quartz, 510
 Quaternary, foraminifers, 186
 nannofossils, 707, 711
 Zonation, 703
 Site 391, 180
 Queyrellin Section, 957
 Radiolarian, Cenozoic, 865
 Hole 391A, 865
 Site 388, 39, 865
 Site 390, 98, 865
 Site 391, 194
 Species list, 867
 Zonation, *Amphirhopalum ypsilon* Zone, 194, 866
 Bekoma bidartensis Zone, 93, 99, 865
 Buryella clinata Zone, 93, 99, 865
 Calocycletta costata Zone, 180, 194, 866, 867
 Cyrtocapsella tetrapera Zone, 180, 194, 866
 Dorcadospyris alata Zone, 180, 194, 866
 Lamprocyrtis haysi Zone, 180
 Lynchnocanoma elongata Zone, 181, 194, 867
 Phormocystis striata Zone, 93
 P. striata striata Zone, 99
 Stichocorys delmontense Zone, 181, 194, 866, 867
 S. wolfii Zone, 194
 Theocampe mongofieri Zone, 865
 Theocotyle cryptocephala cryptocephala Zone, 865
 Thyrsocystis triacantha Zone, 865
 Radiometric time scale, 12
 ages, 735
 Cretaceous stratigraphic units, 738
 Red limestone, see also: Limestone, Red
 Redox conditions, cause of diagenetic migration of
 elements, 581
 Reef banks, 541
 Reef-lagoon facies, Blake Escarpment, 70, 337
 Reflection profiles correlated with drilling, Site 388, 41
 Site 390, 103
 Site 391, 198
 Site 392, 355
 Reflector A, 541
 C, 537, 539
 M, 529
 Blake Plateau, 540
 Regression, Coniacian, 942
 Responsibilities for authorship, 11
Reticulofenestra umbilica Zone, 782, 807
 Sphenolithus furcatulithoides Zone, 703
Retriolpites georgensis Zone, defined, 892
Rhabdosphaera inflata Subzone, 782
 Rhodochrosite, 581
 Ridge and swale topography, 5
 origin of, 23
 Rifting, Atlantic margin, 941
 Rills, 531
 Rim complexes, 541
Rocella gelida, new combination, Incertae Sedis, 788
 schraderi, new species, Incertae sedis, 788
 Saint Crepin Section, 956
 Salisbury Embayment, 941
 Salvadorian Stage, defined, 735
 Sandy limestone, Site 391, 463
 Sapropelitic black shale, 597
Schackoina cabri Zone, 352
 -*Globigerinelloides algerianus* Zones (LC10-11), 353
 Schooner Cays, 424
 oolitic beds, 428
 Scotian shelf, 673, 674, 676
 Jurassic biozonation, 192
 magnetic profiles, 24
 Sediment accumulation rates, Blake-Bahama Basin, 202
 Hole 386, 504
 Hole 387, 504

- Hole 387, 504
- Site 382, 504
- Site 388, 33, 41
- Site 390, 103
- Site 391, 170, 194
- Site 392, 353
- Upper Jurassic red limestone, 503
- Sediment classification system, 14
 - transport across Blake Plateau, 85
- Sedimentary environment, western North Atlantic, 958, 971
- Sedimentary hiatus, Upper Cretaceous, 504
- Sedimentary history, Blake Nose, 85
- Sedimentary structures, fenestral limestone, 433
 - oolitic limestone, 421
 - Site 390, 77
 - skelmoldic limestone, 414
- Sedimentary units, western North Atlantic, 971
- Sedimentation rates, western North Atlantic, 959
- Sedimentologic analyses, Site 388, 482
 - Site 390, 484
 - Site 391, 488
 - Site 392, 489
- Sedimentological features, Blake-Bahama Basin, 949
- Sediments, Hole 391C, 523
 - Site 388, 482
 - Site 390, 483
 - Site 391, 486
- Seismic reflection profiles, 153
 - evidence from, 23
 - stratigraphic correlation of, 27
- Seismic survey and underway data, 11
- Sheet cracks, 424
- Shelf-basin fractionation, 745
- Siderite concretions, 32
- Silicoflagellates, *Corbisema? xenica*, new species, 816
 - toxema*, new species, 815
 - Dictyochoa deflandrei lobata*, new subspecies, 785
 - Distephanus crux loeblichii*, new subspecies, 817
 - ?acanthicus*, new species, 816
 - triotus*, new species, 818
 - Mesocena diodon nodosa*, new subspecies, 818
 - ?connudata*, new species, 786
 - ovata*, new species, 786
 - Naviculopsis eobiapiculata*, new species, 787
 - foliacea tumida*, new subspecies, 820
 - obtusarca*, new species, 865
- Site 382, 775
- Site 384, 775
- Site 385, 778
- Site 386, 780
- Site 387, 782
- taxonomy, 783, 814
- zonation, 783, 812
 - Corbisema triacantha* Zone, 809, 811, 812
 - Dictyochoa hexacantha* Zone, 783, 787
 - spinosa* Subzone, 779, 814
 - spinosa* Zone, 783
 - Naviculopsis biapiculata* Zone, 779, 783
 - constricta* Zone, 776, 783, 786
 - foliacea* Zone, 783, 787, 812, 814
 - robusta* Subzone, 814
- Site 367, correlation with Site 391, 899
 - Jurassic, 902
- Site 382, nannofossils, 775
 - sediment accumulation rates, 504
 - silicoflagellates, 775
- Site 384, nannofossils, 775
 - silicoflagellates, 775
- Site 385, nannofossils, 778
 - silicoflagellates, 778
- Site 386, nannofossils, 780
 - silicoflagellates, 780
- Site 387, nannofossils, 782
 - silicoflagellates, 782
- Site 388, background and objectives, 23
 - benthic forams, 36
 - biostratigraphy, 34, 657
 - CCD, effect on microfossil assemblage, 657
 - foraminifers, 35
 - gas clathrates, 33
 - geochemistry, 33
 - interstitial water, 34
 - lithology, 30
 - lower continental rise hills, 23
 - micrograin size analyses of sediments, 482
 - mineralogy of sediments, 482
 - mystery fossils, 40
 - nannofossils, 37, 707, 807
 - operations, 28
 - organic chemical analyses, 33, 605, 645
 - physical properties, 34
 - radiolarians, 39, 865
 - reflection profiles correlated with drilling results, 41
 - sediment accumulation rate, 33, 41
 - sedimentologic analyses, 482
 - sediments, 482
 - siderite concretion, 32
 - summary and conclusions, 42
 - X-ray mineralogy, 515
- Site 388 to Site 389, underway data, 547
- Site 389, biostratigraphy, 657
 - nannofossils, 707, 807
 - operations, 70
- Site 390, biostratigraphy, 657
 - coarse fraction of sediments, 484
 - Cretaceous, 93
 - ostracodes, 921
 - depositional environment, 99
 - geochemistry, 86
 - interstitial water, 86
 - micrograin size analyses, 485
 - mineralogy, 485
 - nannofossils, 707, 739, 807
 - operations, 73
 - organic geochemical analyses, 606
 - physical properties, 86
 - radiolarians, 98, 865
 - reflection profiles correlated with drilling results, 103
 - sediment accumulation rates, 103
 - sedimentary structures, 77
 - sedimentologic analyses, 484
 - sediments, 483
 - subsidence rates, 103
 - Tertiary, 93
 - Upper Cretaceous, stratigraphic hiatus, 504
 - X-ray mineralogy, 516
 - Zoophycos, 77
- Site 390 to Site 391, underway data, 554
- Site 391, background and objectives, 153
 - biostratigraphy, 179, 660
 - Blake-Bahama Basin, 153
 - chlorins, 669
 - coarse fraction of sediments, 488

- Cretaceous-Upper Jurassic, 181
- depositional environment, 181
- diagenesis, carbonate sediments, 478
- environments of deposition, Cretaceous-Jurassic limestone, 465
- foraminifers, 181
- geochemistry, 171
- gravity transport, 162, 201
- Green River shale compared to, 636
- hydrocarbon generation, 618
- interstitial water chemistry, 171
- lithology, 160
- micrograin size analyses, 488
- mineralogy, 488
- nannofossils, 711, 741, 768, 808
- organic carbon content, 626
- organic chemical analyses, 171, 606, 617, 625, 646
- petroleum-bearing potential, black claystones, 171, 617, 625
- physical properties, 171
- radiolarians, 194
- reflection profiles correlated with drilling results, 198
- sediment accumulation rates, 170, 194
- sedimentologic analyses, 488
- sediments, 486
- stratigraphic hiatus, Miocene/Cretaceous, 203, 625, 630
- summary and conclusions, 201
- thermal history, 643
- X-ray mineralogy, 517
- Site 391 to 392, underway data, 554
- Site 392, background and objectives, 337
 - biostratigraphy, 351, 657
 - depositional environment, 346, 353
 - diagenesis, 356
 - foraminifers, 351
 - geochemistry, 349
 - lithology, 339
 - micrograin size analyses, 490
 - microkarst formation, 349
 - mineralogy, 489
 - nannofossils, 353, 707, 810
 - oolite, 343
 - operations, 337
 - ostracodes, 921
 - physical properties, 349
 - reflection profiles correlation with drilling results, 355
 - sediment accumulation rates, 353
 - sedimentologic analyses, 489
 - summary and conclusions, 356
 - X-ray mineralogy, 518
- Site 392 to Norfolk, underway data, 554
- Site 393, background and objectives, 395
 - lithology, 397
 - nannofossils, 711
 - objectives, 395
 - operations, 395
 - summary and conclusions, 398
- Site 394, background and objectives, 395
 - lithology, 397
 - nannofossils, 711, 810
 - objectives, 395
 - operations, 396
 - X-ray mineralogy, 518
- Skelmoldic limestone, 345, 414, 489
 - depositional environment of, 416, 419
 - diagenetic history, 417
 - lithology, 414
 - sedimentary structures in, 414
 - See also:* limestone, skelmoldic
- Slump deformation hypothesis, lower continental rise hills, 44
- Slumping, 958
- Sourelieu Section, 955
 - compared to Leg 44 sediments, 541
- South Jersey high, 943
- Sphaeroidinellopsis subdehiscens* Zone (N 13), 180
- Sphenolithus belemnos* Zone (NN3), 712, 810
 - ciperoensis* Zone, 781, 788, 810
 - furcatolithoides* Zone (NP16), 93, 98, 703, 708, 709
 - defined, 703
 - heteromorphus* Zone (NN 5), 712, 713, 809, 810, 814
- Sponges, 811
- Sporomorphs*, Clavatipollenites Zone, defined, 892
 - Druggidium deflandrei/Ephedripites multicoastatus* zones, 894
 - Ephedripites multicoastatus* Zone, defined, 892
 - Ephedripites multicoastatus-Retitricolpites georgensis* Zone, 890
 - Psilatricolporites* Subzone, 890, 892
 - Retitricolpites georgensis* Zone, defined, 892
 - Tricolpites minutus* Subzone, 890, 892
- Stichocorys delmontense* Zone, 181, 194, 866, 867
- wolfii* Zone, 194
- Stratigraphic palynology, 888
- Stratigraphic analyses, Miocene/Cretaceous, 625
 - correlations, seismic reflectors, 27
 - hiatuses, 958
 - Albian-Campanian, 106, 490, 492, 707
 - Barremian, 953
 - Cretaceous-Paleocene boundary, 106
 - Lower Cretaceous, 98
 - Maestrichtian, 96
 - Maestrichtian-Paleocene, 492
 - Miocene-Cretaceous, 203, 630
 - Paleocene-Eocene boundary, 106
 - Upper Cretaceous, 504
- Strontium concentration, carbonates, 557, 559
- Subaerial weathering, 7
- Subalpine Zone, 953
- Submarine gravity-flow deposits, 8
- Subsidence, western North Atlantic, 959
 - Blake Nose, 951
 - Site 390, 103
 - Sites 389 and 390, 106
- Subtilisphaera perlucida* Subzone, 890, 892
- Systematic descriptions, Calcisphaerulidae, 912
 - nannofossils, 746
 - silicoflagellates, 783, 814
- Tectonic uplift, Cuba, 202
- Tenerifian Stage, defined, 738
- Tertiary, benthic foraminifers, 36
 - biostratigraphy of, 943
- Tertiary-Cretaceous boundary, nannofossils, 93, 98, 710
- Tethys Current, 743
- Tetralithus trifidus* Zone, 93, 98, 353, 710, 810
- Tetrapyrrole pigments, depth-age profile of, 639
 - geochemistry of, 639
 - mass spectrometric analyses of, 639
- Thecampe mongolfieri* Zone, 865
- Thecotyle cryptocephala cryptocephala* Zone, 865
- Thermal analysis/pyrolysis, 635
- Thermal history, Site 391, 643

- Thermal immaturity, evidence for, 626
- Thermal maturation, 645
- Thermohaline circulation, 27
- Thin-layer chromatography, 593, 621
- Thyrsocypris triacantha* Zone, 93, 99, 865
- Ticcinella breggiensis* Zone (LC 17), 352, 664, 670
 - primula* Zone (LC15-16), 93, 97, 353, 664, 669, 671
 - Ticcinella breggiensis* Zones, 673, 676
- Titanium, concentration in sediments, 576
 - red limestone, 505
 - variegated and red claystone
- Toco and Cuhe formations, 672
- Tongue of the Ocean, marine sediment source, 181
- Trace elements in carbonate, 557
- Transgression, Albian-Turonian, 942
- Tribrachiatius orthostylus* Zone, 783, 808, 814
- Tricolpites minutus* Subzone, 890, 892
- Triquetrorhabdulus carinatus* Zone, (NN1), 181, 712
- Trithrodinium suspectum/Complexipollis* zones, 893
- Tubodiscus jurapelagicus* Zone, 714
- Turbidites deposition, Pleistocene, 33
- Turbidites, 971
 - Blake-Bahama Basin, 207
 - Pleistocene, 33
 - Upper Miocene, 153, 204
- Turbidity current, 463, 485, 489, 493
- Unconformity, Campanian-Albian, 541
- Underway data, collection and presentation, 547
 - Norfolk to Site 388, 547
 - Site 388 to Site 389, 547
 - Site 390 to Site 391, 554
 - Site 391 to Site 392, 554
 - Site 392 to Norfolk, 554
- Upper Cenozoic, foraminifers, 95
- Upper Cretaceous, foraminifers, 96, 352
 - lithostratigraphy and biostratigraphy, 942
 - nannofossils, 98, 710
 - stratigraphic hiatus, 504
- Upper Jurassic, Calcisphaerulidae, 911
 - limestone, 154
 - nannofossil zonation, 710, 762
- Upper Miocene, turbidites, 153, 201, 204
 - Site 391, 180
- Urgonian Facies, 953
- Utatur formation, 676
- UV-visible spectra; data, pigment fractions, 639
- variegated and red claystones, aluminum content, 507
 - magnesium content, 508
 - manganese content, 506
 - origin of, 515
 - potassium content, 508
 - silica content, 508
 - titanium content, 506
- Velocity analyses, 533
- Verrill Canyon Formation, 676
- Vitrinite, 585, 602
 - reflectance of, 599, 617, 618, 621, 629
- Watznaueria communis* Zone, 767
- Western Atlantic, continental margin, 23
 - depositional environment, 958, 971
 - marine bottom topography, 958
 - rate of subsidence, 959
 - sedimentary environment, 958
 - sedimentary units, 971
 - sedimentation rates, 959
- Western North Atlantic and north Tethyan margins,
 - comparison of, 957
- X-ray mineralogy, methods of, 515
 - Site 388, 515
 - Site 390, 516
 - Site 391, 517
 - Site 392, 518
 - Site 394, 518
- Zeolites, 511
- Zinc, concentration in sediments, 579
- Zoophycos, 77