

INDEX

- Aaiun basin, 77, 80, 735, 779, 790, 808, 810
 deltaic continental to lagoonal facies, 782
Early Cretaceous deltaic complex from, 34
Late Cretaceous transgression, 783
 sedimentary sequence, 559
Aaiun-Tarfaya coastal basin, 19, 568, 811
 Mid-Cretaceous organic-rich shales, 804
 subaerial deltaic muds, 78
Acoustic horizons, 456
 Site 397, 448
 velocity, Site 397, 41
African climatic history reconstructed from coarse fraction analysis, 662
African margin off Cape Bojador, sequence of events in Neogene history of, 36
 petroleum potential, 78
 rapid base-of-slope-fill from, 611
Agadir Basin, 261
Agadir-Essaouira Region, correlated with offshore pro-delta facies, Site 397, 261
Age of sediments, Lower Cretaceous, 226
 Canary Islands, volcanic apron at southeastern foot, 794
Air-fall ash deposits from Gran Canaria Island, 32
Air-Hoggar Mountains, 802
Albian to Turonian transgression, 19
Albian to Upper Cretaceous sedimentation rate, 488
Algal type organic matter, 506, 511
Alkalic microgabbros, 710
Alpine orogeny, High Atlas, 785
 proximal flysch sequences, 792
Alps, Lower Cretaceous faunas, 582
Alteration of volcanoclastic sediments, Site 397, 746
Alternating-field demagnetization, 464
Amaurolithus primus Zone, 380
Ammonites, 283
 biostratigraphy, 68
 macrofossils, 30, 34
 neocomites, 283
Analcime, 746
Antarctic bottom water, 665, 790
 convergence, 84
 glaciation, Eocene/Oligocene, 83
 late Miocene, 36
 effect on sediment transport, 661
ice cap, expansion of, 36
 onset of glaciation, 785
ice caps, deglaciation, 64
ice sheet, Pliocene, reduction in size during Pliocene, isotopic evidence, 443
 late Miocene extent, 664
Antarctica, glacial expansion in Miocene, 455
Anti-Atlas Mountains, 80, 567
Anti-Atlas/South Atlas Fault (Cape Juby), 77
Aptian/Albian, global increase of subsidence rates, 811
 Moroccan coastal basins, 811
Arctic glaciation, 665
 late Pleistocene, 662
Arctic Ocean, early Pleistocene, constant conditions in, 662
 Pleistocene ice conditions, diatom evidence, 479
Ash layers, burrowing in, 704
 sideromelane shards, 711
 Site 397, 703
 composition, 707
Atlantic deep-water, effects on benthic communities, 735
 circulation, Quaternary changes in, 419
 changes, surface-water regime changes correlated with, 735
Atlas Mountains, 802
 chlorite source, 643
 source of smectite, 616
 tectonics, 663
Attapulgite, 600
Authigenic phyllosilicates, 748
 minerals, age relationship between, 748
 silica, 751
Authorship, responsibilities for, 5
Autochthonous sequence, Site 397, Neogene sediments, 647
Background and objectives, Site 397, 19
Bahama Basin and Bermuda Rise, metalloporphyrins, 550
Baie du Levrier, sediments, 736
Baltimore Canyon Trough, 810
 geothermal gradient, 805
 subsidence rates, 809
Basalt, Canary Islands, shield building, 714
 fragments, chemical analysis of, 700
Bavarian Limestone Alps, foraminiferal genus, *Favusella*, 273
Benthic foraminifers, biostratigraphy, 60, 422
 Central North Pacific, 60
 displacement, 422
 Early Cretaceous, 63
 faunal fluctuations, *Meteor* cores, 731
 grain-by-grain transport, 422
 isotopic curves, *Meteor* cores, 731
 Meteor cores, 731
 Neogene and Quaternary, 60
 oxygen and carbon isotope stratigraphy, 433
 paleobathymetry, 62
 Site 367, 60
 369, 60
 397, 419
Benthic/planktonic ratio, foraminifers, 653
 ratios, foraminifers, 666
Betancuria Massif, Fuerteventura Island, 713, 794

- Biochronology, foraminifers, 327
 Biodegradation of organic matter, 503
 Biogenic, methane, 531, 532
 - productivity effects, upwelling, 611
 - sedimentation in region of Cape Bojador, 35
 Biogenous volcanic sandstones of the Emperor Seamounts, 748
 Biological methane, 804
 Biostratigraphic evidence for submarine erosion, 57
 Biostratigraphy, ammonites, 68
 - and magnetostratigraphy, Site 397, 85
 - benthic foraminifers, 60, 422
 - foraminifers, 67
 - Lower Cretaceous, Hole 397A, 581
 - Site 397, 561
 - magnetostratigraphic evidence, 68
 - nannofossils, 67
 - sedimentological evidence, 68
 - seismic evidence, 68
 - Site 397, 51
 - foraminifers, 257- Biotite diagenesis, 581
 Bioturbation, 35, 560, 607, 612, 781, 791
 - effects on climatic record, 433
 - structures, 28
- Black Sea, Hole 381, 537
 - hydrocarbon migration, 535
 - metalloporphyrins, 550
 - N*-methyl ketones, 542
 - Sites 379-381, 531
- Blake Plateau Basin, 810
 Blake-Bahama Basin, nannofossils, Leg 1, 227
 Boron, concentration in sediments, optical emission spectrophotometry analyses, 553
 - geochemistry, Miocene/Pliocene boundary, 553
 - in sediments, 553
- Bou Reg Reg section of Morocco, 453
 Boudinage-structure, siderite layers, 560
 Boudinage-type features, 779
 Brunhes Normal Epoch, 472, 479
 - paleomagnetic Epoch, 62, 69, 405, 422, 457, 464, 470, 475, 672
 - to Epoch 7, 86
- Brunhes/Matuyama boundary, 472
 Bulk physical properties, Site 397, 44
 - properties of sediments, Site 397, 45
- Burrowing in ash layers, 704
 Calabria, Le Castella section of, Pliocene/Pleistocene boundary stratotype at, 456
 Calcareous components, diagenetic potential of, 87
Calcicalathina oblongata Zone, 66, 226
 Calcite, cementation by, 747
 - compensation depth, 487
- Calcium carbonate dissolution, 31, 32
 - early Miocene, Equatorial oceans, 797
 - Quaternary, Site 397, 665
 - upper Miocene, Site 369, 665
 - 397, 665
- California Borderland, flysch sequences from, 617
 Canary Current, 36, 381, 391, 643, 665, 797
 Canary Island Archipelago, 779
 - subaerial volcanic stage of, 811
 - volcanic uplift, Neogene, 713
 - area, heat flow anomaly in, 805
 - channel, 807
 - paleodepth, 785
 - turbidites in, 84
 - turbidity currents, 715
 - ridge, uplift of, 790
 - shelf, source of sediments, Hole 397, 31
 - volcanic debris, 613
 - volcanism, 801, 811
 - effects, thermal gradient, 70
 - evolution of, 794
 - middle Miocene, 86, 87
 - Neogene evolution, 703
 - volcanoes, mid-Miocene uplift of, 792
- Canary Islands, 775
 - debris flows from, 35
 - early evolution of, 715
 - early to middle Miocene, uplift and shield stage, 794
 - marine terraces, 66
 - potassium-argon age determinations of volcanic samples from, 666
 - shield building, basalt, 714
 - Site 138, west of the, 790
 - uplift, contribution to smectite concentration at Site 397, 600
 - uplift event, late-early to middle Miocene, 37
 - volcanic apron at southeastern foot, age of sediments, 794
 - volcanism, 701
 - volcanic center, effect on geothermal gradients, Site 397, 508
 - volcaniclastic flows from, 616
 - volcanism, Miocene, 615
- Cape Bojador, 19, 221, 375, 419
 - biogenic sedimentation in region, 35
 - continental margin, changes in interpretation, 19
 - deltaic sequence, 568
 - Early Cretaceous paleoenvironment, sedimentary history, 559
 - evolution of, 811
 - Northwestern Africa, geologic history of, 773
 - paleobathymetry, Mesozoic/Cenozoic, 807
 - subsidence, 568, 807, 809
 - continental shelf of, 779
 - depositional sequences off, 485
 - lower slope/upper rise region off, 67
 - margin, Wealden sequence at the, 810
 - marginal basin, 779, 804, 807
 - removal of pre-Miocene sediment, 785
 - Neogene sediments in region of, 298
 - northwest African continental margin off, 615, 617, 785
 - slope, seismic interpretation, 491

- Cape Cod, Georges Bank Basin off, 811
 Cape Juby, 779
 to Cape Blanc, continental margin, 783
 Cape Verde Basin, Site 367, 221
 Site 370, 221
 Islands, Lower Cretaceous pelagic carbonates and shales, 794
 Rise, contourites, Site 368, 793
 early Miocene uplift of, 37
 uplift of, 790
 Cape Verde-Madeira Abyssal Plain, 19
 Capo Rossello, 384
 Carbon isotope fluctuations, Pleistocene, 443
 Carbon preference indexes, 516
 Carbon-carbonate analyses, 13
 Carbon-isotope record, *Meteor* cores, 735
 results, *Meteor* cores, foraminifers, 728
 Carbon-13 values, forestation effects on, 398, 442
 Methane, 533, 809
 Site 397, 86, 398, 508
 Methane, 804
 Carbon-14 record, northwestern Sahara Desert, climatic fluctuations, 736
 Carbonate bomb measurements, 11
 cement, volcanic sandstones, 713
 compensation depth, 34, 64, 671, 675
 content, Site 397, late Neogene, 671
 curve, 680
 curves, foraminifer, 679
 curves, stable isotopic ratios correlated with, 671
 diagenesis, 86
 values, upper Miocene, Site 397, 672
 Carbonification, katagenesis stage, 587
 Cariaco Trench, particulate organic matter concentrations in waters of, 504
 Site 147, 531
 Cementation by calcite, 747
 silica, 747
 smectite, 747
 zeolite, 747
 Cenomanian to Turonian sedimentation rates, 568
 transgression, 568, 809
 Cenozoic abundance, nannofossils, 296
 preservation, nannofossils, 296
 Central North Pacific, benthic foraminifers, 60
Ceratolithus acutus Zone, 380
rugosus Zone, 380
tricorniculatus and *Ceratolithus rugosus* zones (NN 12-13), 293
tricorniculatus Zone (NN 12) to *Discoaster brouweri* Zone (NN 18), 64
 Chabazite, 746
 Chalk diagenesis, Site 167, equatorial Pacific Ocean, 674
 Site 397, 674
 Change in *iC5/nC5* ratio at stratigraphic hiatus, lower Miocene/Hauterivian, 534
 Changes in interpretation of Cape Bojador continental margin, 19
 Chemical analyses, Site 397, sediments, 49
 analysis of basalt fragments, 700
 overprinting, Sites 379 and 380, paleomagnetic measurements, 464
 Chlorins, 548
 Chlorite, 643
 source, Atlas Mountains, 643
 Chloroform extracts, 525
 Chlorophyll, diagenesis of, 547
 Site 397, 547
 Chondrites, 608
 Circum-Antarctic bottom-water circulation, 83
 current, 785
 Clay fraction analysis, Site 397, 643
 mineralogy, Site 397, 595
 minerals, 560
 X-ray diffraction results, 32
 studies of, Site 397, 595
 Clays, Lower Cretaceous, geochemistry of, 583
 Climatic fluctuations, oxygen-isotopic record evidence, late Pleistocene, 797
 change and desertification in Africa, deep sea record of, 802
 cycles identified with, oxygen-isotope data, 727
 fluctuations, carbon-14 record, northwestern Sahara Desert, 736
 correspondence to North American glacial cycles, late Pleistocene, 797
 Holocene, 727
 late Pleistocene, 727, 733
 history, late Neogene, 433
 northwestern Sahara coastal margin, 737
 Clinoptilolite, 746, 748, 751
 Clinopyroxene, volcaniclastic sandstones, igneous components, 710
 Coarse fraction analyses, glauconite, Site 397, 658
 late Miocene, Site 369, 663
 Meteor cores, 734
 pyrite, Site 397, 658
 relic particles, Site 397, 658
 terrigenous material, Site 397, 658
 volcanic material, Site 397, 658
 analysis, phosphate grains, Site 397, 660
 Quaternary/Pliocene boundary, 660
 Site 397, 641, 647
 volcanism, 666
 results, summary of, Site 397, 30
 Coarse frosted-quartz evidence, dune formation, 32, 84
 Compaction, 768
 physical properties, sediments, 761
 sedimentary basin, 761
 Computer modeling, Site 397, 765
 GRAPE measurements, physical properties compared with, 766
 pore pressure, 765
 thermodynamic parameters, 765
 Continental climate and deep-sea record, Site 397, 802
 margin, Cape Juby to Cape Blanc, 783
 cores, oxygen-isotope curves, 728
 destruction, possible causes of, 785
 of the eastern North Atlantic, terrigenous input to, 736

- rise, defined for continental margin of northwest Africa, 447
 near Cape Bojador, 19
 western African, 727
 shelf off Cape Bojador, 779
 slope of northwestern Africa, lower and middle Miocene, sandy sediment, grain morphology, 683
 grain-size, 683
 Contourites, Site 138, 793
 Site 368, Cape Verde Rise, 793
 Contribution to smectite concentration at Site 397, Canary Islands, uplift, 600
 Core disturbance, Leg 47A, 7
 Cretaceous black shales, 541
 nannofossils, 52
 sedimentation rates, 65
 (post Hauterivian), Site 369, sediments, 782
 mollusks, 64
 Cretaceous-Paleogene Carpathian flysch, 84
Cretarhabdus loriei Zone, 226
Cruciellipsis cuvilliéri Zone, 53
 Crystalline basalt, volcanoclastic sandstones, igneous rock fragments, 710
 Cyclic sediment transport, lower and middle Miocene, 695
 Cycling of organic matter in the world's oceans, 503
 C1 to C5 hydrocarbon analysis, Leg 42B, 47
 hydrocarbon analysis, organic geochemistry, 47
 C1 to C7 hydrocarbons, 536
 Site 397, 531
 C2 to C4 hydrocarbons diagnostic of petroleum formation, 531
 C2 to C5 hydrocarbons, 531, 534, 536
 C6 to C7 hydrocarbons, 536
 Debris flows, 606, 782
 Quaternary, passive continental margin of northern Africa, slumps, 614
 from Canary Islands, 35
 northwest African margin, 35
 Decollement zones, sediment structures, 781
 Deep-current erosion causing slope-rise destruction, 34
 Deep-sea record of climatic change and desertification in Africa, 802
 Deglaciation, Antarctic ice caps, 64
 Dehydroabietic acid, 542
 Deltaic continental to lagoonal facies, Aaiun Basin, 782
 sediments, Lower Cretaceous, source area in, 567
 sequence, Cape Bojador continental margin, 568
 Demerara Rise, Site 144, 566
 Density, physical properties, fluids, 761
 Depositional environment, Early Cretaceous sedimentary basin, 585
 unstable salinity regime, 585
 Site 397, Lower Cretaceous foraminifers, 257
 environments and processes, Site 397, 33
 sequences off Cape Bojador, 485
 Desert quartz number, 658
 numbers, 641, 659, 661, 662
 Diagenesis, biotite, 581
 Lower Cretaceous sediments, Site 397, 567
 organic matter, 775
 siliceous organisms, 742
 organic matter and hydrocarbon gases, 86
 pelagic carbonate sediments, 448
 silica, 750
 of chlorophyll, 547
 Diagenetic potential of calcareous components, 87
 Diatom evidence, Pleistocene ice conditions, Arctic Ocean, 479
 Diatoms, Site 397, 653
Diloma, new nannofossil genus, 228
 placinum, new nannofossil species, 228
 primitiva, new nannofossil combination, 228
Discoaster asymmetricus Zone (NN 14), 294
 berggrenii Subzone, 380
 berggrenii Zone, 379
 brouweri Zone (NN 18), 51, 294, 381
 calcaris Zone (NN 10), 52, 293
 druggii Zone (NN 2), 64, 221, 289, 291
 exilis Zone (NN 6), 52, 293
 hamatus Zone (NN 9), 293, 377
 to *Discoaster quinqueramus* Zone (NN 11), 64
 298
 kugleri Zone (NN 7), 52, 293
 neohamatus Zone, 377
 pentaradiatus Zone (NN 17), 51, 294, 381
 quinqueramus Zone (NN 11), 293
 surculus Zone, 51, 294, 381
 tamalis Zone, 384
 Dispersed organic matter (DOM), composition, 584
 methods of study, 584, 594
 transformation, 584
 Displacement, benthic foraminifers, 422
 of Samlat Formation rocks by slumping, 20
 Dissolution at depth, foraminifers, Site 397, 56
 index, foraminifers, 672, 679
 measure, foraminifers, 672, 679
 ratio, 673
 Distal delta facies accumulation, Lower Cretaceous sediments, Site 397, 260
 prodelta facies of Early Cretaceous prograding delta, 782
 muds, 808
 turbidites, 509, 794, 806
 Downhole temperature, 505
 Downhole temperature measurements, Site 397, 69
 Downslope displacement, sediment transport, 737
 sedimentary processes, 55
 Drilling information, Sites 369, 491
 operations, Hole 397, 21
 397A, 23
 results, Seismic reflection profiles correlated with, 70
 Drusbergschichten (Barremian) of Vorarlberg (western Austria), 273
 Dune formation, coarse frosted-quartz evidence, 32, 84
 Dust-derived sediments, Sahel lateritic belt, Sahara Desert, 803
 Early evolution of the Canary Islands, 715
 Early to middle Miocene, initial sedimentation phase, 36
 uplift and shield stage, Canary Islands, 794
 Early Cretaceous, benthic foraminifers, 60, 63
 deltaic complex from Aaiun Basin, 34
 Hole 397A, mollusks, 283

- New Foundland continental margin, 567
 Nova Scotian continental margin, 567
 paleodepths, Site 397, 809
 paleoenvironment, 561
 sedimentary history, Cape Bojador continental margin, 559
 paleomagnetic evidence, sedimentation rates, 475
 prograding delta, distal prodelta facies of, 782
 sedimentary basin, depositional environment, 585
 unstable salinity regime, depositional environment, 587
 sedimentation rates, 782
 stratigraphic hiatus, 60
 subsidence rates, 809
 Wealden facies, 568
 Wealden-type delta foreset beds, 807
 Early Miocene, along African continental margin south of Canary Islands, upwelling, 798
 equatorial oceans, calcium carbonate dissolution, 797
 porcelaneous mudstones, 751
 sedimentation rate, 612, 764
 uplift of Cape Verde Rise, 37
 upwelling, 751
 Early Neogene displaced sediment regime, 790
 Early Pleistocene, constant conditions in Arctic Ocean, 662
 Earthquake activity, Miocene, related to Paleogene-Neogene organic phases in Atlas Mountains, 615
 Earthquakes, 801
 Eastern boundary currents, 788
 Eastern Antarctic ice cap, middle Miocene upwelling with buildup of, 798
 Echinoderms, Site 397, 653
 Electron microscopy studies, sediment diagenesis, 742
Emiliania huxleyi Zone (NN 21), 51, 289, 296
 Emperor Seamounts, biogenous volcanic sandstones of, 748
 Environmental evolution, late Neogene, 447
 Eocene through Oligocene paleoclimates, 785
 Eocene/Paleocene, stratigraphic hiatus, 485
 seismic reflection profile evidence, 488
 Eolian material, Site 397, 695
 Eolian origin of lipids, 541, 543
 Eolian sand turbidites, Site 139, 793
 Site 397, 644
 Eolian transport of volcanic material, Site 369, 666
 Equatorial oceans, calcium carbonate dissolution in early Miocene, 797
 Equatorial Pacific Ocean, thermohaline bottom currents, 783
 chalk diagenesis, Site 167, 674
 Erionite, zeolites, 747
 Essaouria Basin, 261
 Euryhaline fauna from Paratethys, 455
 Eustatic sea-level rise, 261
 Euxinification of the Mediterranean Sea, 451
 Even/odd carbon preference, organic matter, 542
 Evolution of Canary Island volcanism, 794
 Cape Bojador continental margin, 811
 passive margins, 617
 Expansion of Antarctic ice cap, 36
 Explanatory notes, 6
 Extract quantity and type, 505
 Facies migration, 568
 Faults, 779
 Faunal fluctuations, *Meteor* cores, benthic foraminifers, 731
 Feldspar, volcaniclastic sandstones, 710
 Ferriferous carbonate bands, sedimentation of, 586
 Fine-grained volcanic debris from Fuerteventura shelf, 32
 Fining-upward facies sequence, 568, 614, 809
 Fish debris, Site 397, 560, 653, 664
 Flaser bedding, 560
 Flow structures, 779
 Fluid and heat flow, 761
 model, 764
 Fluid flow, 761
 Fluids, density, physical properties, 761
 thermal properties, 761
 viscosity, physical properties, 761
 Fluvial sediments offshore, Sequia El Hamra Wadi, 616
 Fluxoturbidites, 792
 Flysch, 84
 sedimentation, 78
 sequence, 794
 sequences, Alpine proximal, 792
 from California Borderland, 617
 Flyschoid section, 774
 sediments, 804
 Fold structures, sedimentary structures, 612, 781
 Foraminifers, benthic/planktonic ratio, 653, 666
 biochronology, 327
 biostratigraphy, 67
 carbon-isotope results, *Meteor* cores, 728
 carbonate curve, 679
 dissolution index, 672, 679
 measure, 672, 679
 Early Cretaceous, 60
Favusella stiftia, new species, 276
 fragmentation of, 648, 658
 Genus, *Favusella*, 273
 Bavarian Limestone Alps, 273
 Drusbergschichten (Barremian) of Vorarlberg (western Austria), 273
 Scotian Shelf of Atlantic Canada, 273
 Site 105, 274
 Site 144, 274
 Site 97, 274
 Lower Cretaceous, biostratigraphy, Site 397, 257
 depositional environment, Site 397, 257
 paleoecology, Site 397, 257
Meteor cores, 731
 Miocene, 59, 317
 stratigraphy, Site 372, 321
 Miocene/Pliocene boundary, 59, 318, 327, 341, 673
 N.12/N.13 zonal boundary, 337
 N.13/N.14 zonal boundary, 321
 N.15/N.16 zonal boundary, 57
 N.21/N.22 zonal boundary, 59
 N.6/N.7 zonal boundary, 59
 N.8/N.7 zonal boundary, 59

- Neogene, sedimentation rates based on, 64
 oxygen-isotope results, *Meteor* cores, 728
 paleotemperature curves based on, 735
 Pliocene, 58
 Pliocene/Pleistocene boundary, 57
 Quaternary, 57
 Site 397, 53
 - dissolution at depth, 56
 - preservation, 57
 solution index, 671
 Tortonian/Messinian boundary, 339
 Zonation, *Globigerina nepenthes* Zone, 58, 338, 339
 - nepenthes/Globorotalia siakensis* Zone, (N.14), 59
 - Globigerinatella insueta/Globigerinoides quadrilobatus trilobus* Zone (N.7), 59
 - Globigerinoides elongatus* Zone (M Pl 5), 58
 - sicanus/Globigerinatella insueta* Zone (N.8), 59
 - Globorotalia acostaensis acostaensis/Globorotalia merotumida* Zone (N.16), 59
 - fohsi* Zone (N.12), 59
 - inflata* Zone (M Pl 6), 58
 - margaritae/Globorotalia puncticulata* concurrent-range zone, 58, 339
 - opima opima* Zone (P.21), 67, 611, 782
 - praefohsi* Zone (N.11), 67, 611, 782
 - puncticulata* Zone (M Pl 2/M Pl 3 zonal boundary), 317
 - truncatulinoides* total range zone, 58
 - Orbulina suturalis/Globorotalia peripheroronda* Zone (N.9), 59
 - Sphaeroidinellopsis acme* Zone, 455
 - seminulina paenedehiscens* Zone, 59, 339
 - subdehiscens* partial range zone, 58
 - subdehiscens/Globigerina druryi* Zone (N.13), 59
 Zone M Pl 3, 55, 339
 - M Pl 4, 55, 57, 456
 - M Pl 4/M Pl 5 boundary, 384
 - M Pl 5, 57, 384
 - M Pl 6, 57
 - N.10, 318, 321, 336, 339, 341
 - N.11, 321, 336
 - to N.14, 339
 - N.12, 62, 337
 - N.13, 55, 326, 337
 - N.13/N.14 boundary, 337
 - N.14, 55, 318, 321, 337
 - N.15, 318, 323
 - N.16, 55, 318, 323, 327, 337, 339, 341
 - N.17, 59, 326
 - N.18, 341
 - N.18/N.19 boundary, 327
 - N.19, 327
 - to N.17, 327
 - N.6, 318, 327
 - N.7, 317, 318, 321, 327, 339
 - N.7/N.8 boundary, 318
 - to N.9, 339
 - N.8/N.9 boundary, 318- N.9, 62, 318, 336
- Zones N.10 to N.13, 55
 - N.11 to N.13, 55, 321
 - N.12 to N.13, 55
 - N.14, 84
 - N.16, 60
 - N.13 and N.14, 55
 - to N.19, 323
 - to N.7, 326
 - N.16, 84
 - N.16/N.17, 327
 - N.17 to N.19, 62
 - to N.20, 323
 - N.17/N.18, 327
 - N.2 to N.14, 55
 - N.5 to N.13, 55
 - to N.6, 59, 62, 67, 782
 - N.6 to N.12, 55
 - to N.21, 323
 - N.7 to N.13, 55
 - N.8 to N.10, 55
 - P.18 to N.13, 55
- Forestation, effects on carbon-13 values, 398, 442
- Fragmentation of foraminifers, 648, 658
- Free-base porphyrins, 548
- Fuerteventura volcanic debris flows, seismic reflection profiles, middle Miocene, 795
 - shelf, fine-grained volcanic debris from, 32
 - Island, 85, 493, 613, 704, 794, 795
 - Betancuria Massif, 713
 - volcanic and sedimentary detritus from, 33
 - components source, Site 397, 714
- Fusinite type organic matter, 514
- Gaada Formation, Late Cretaceous black shales, 804
- Gabon-Angola Basin, 804
- Gas chromatography, 531, 541
 - analyses, tetrapyrrole pigment content, 547
- Gauss Paleomagnetic Epoch, 69, 383, 443, 456, 457, 471, 479
- Gauss/Matuyama boundary, 391, 443
- Generation of hydrocarbons, 585
- Geochemical measurements, 13
- Geochemistry, Miocene/Pliocene boundary, boron, 553
 - of Lower Cretaceous clays, 583
 - Site 397, 46
- Geologic history of Cape Bojador continental margin, 773
 - history and evolution of passive continental margin in the eastern North Atlantic during the Mesozoic, 257
 - history of Northwestern African continental margin, 775
- Georges Bank Basin off Cape Cod, 811
- Geothermal gradient, 518, 805
 - Baltimore Canyon Trough, 805
 - Site 397, 508, 512
 - gradients, 766
 - Site 397, 764
 - heat flux, 761
- Gephyrocapsa oceanica* Zone (NN 20), 51, 289, 296

- Gilbert Paleomagnetic Epoch, 69, 384, 391
 Epoch/Epoch 5 boundary, 69, 341
 Glacial expansion in Antarctica, Miocene, 455
 tills in, Iceland, 443
 Glacioeustatic regression, Northern Hemisphere ice-sheet development, 85
 Glass shards, constituents of volcaniclastic sandstones, 699
 Glauconite, Site 397, coarse fraction analyses, 658
 Global increase of subsidence rates, Aptian/Albian, 811
Globigerina nepenthes Zone, 58, 338, 339
nepenthes/Globorotalia siakensis Zone (N.14), 59
Globigerinatella insueta/Globigerinoides quadrilobatus trilobus Zone (N.7), 59
Globigerinoides elongatus Zone (M Pl 5), 58
sicanus/Globigerinatella insueta Zone (N.8), 59
Globorotalia acostaensis acostaensis/Globorotalia merotumida Zone (N.16), 59
fohsi Zone (N.12), 59
inflata Zone (M Pl 6), 58
margaritae/Globorotalia puncticulata concurrent-range zone, 58, 339
opima opima Zone, 21, 67, 611, 782
praefohsi Zone (N.11), 59
puncticulata Zone (M Pl 2/M Pl 3 zonal boundary), 317
truncatulinoides total range zone, 58
 Gmelinite, 747
 Grain morphology, sandy sediments, continental slope of northwestern Africa, lower and middle Miocene, 683
 Grain-by-grain transport, benthic foraminifers, 422
 Grain-size, curve, 680
 ratio, 673
 sandy sediments, continental slope off northwestern Africa, lower and middle Miocene, 683
 Site 397, late Neogene, 671
 Gran Canaria and Fuerteventura islands, oceanic basement between, seismic reflection evidence, 497
 Gran Canaria Island, 704, 779, 794, 809
 air-fall ash deposits from, 32
 magma evolution, 796
 marine terraces on, 85
 middle Miocene volcanic apron of, 497
 rhyolitic eruption on, 85
 submarine shield of, 796
 volcanic components source, Site 397, 714
 Grand Banks, 810
 GRAPE measurements, 15
 physical properties compared with, computer modeling, Site 397, 766
 Greenland, separation from Europe, 785
 Guadalupe delta, prodelta sediments, 564
 Gulf of Mexico, isotopic curves, 735
 Hauterivian, Site 397, sedimentation rate, 564
 mudstones, 603, 774
 sediments, 86
 shales, 523
 Hauterivian, oceanic circulation, 568
 Heat flow, 761, 768
 anomaly in Canary Island area, 805
 measurements, 15
Helicopontosphaera ampliaperta Zone (NN 4), 52, 293, 298, 299
sellii Zone, 295
 Hierro Island, 794
 High Atlas, Alpine orogeny, 785
 Hoggar Mountains, 802
 Hole 381, Black Sea, 537
 Hole 397, Canary Island shelf, source of sediments, 31
 drilling operations, 21
 Hole 397A, biostratigraphy, Lower Cretaceous, 581
 drilling operations, 23
 hydrocarbon generation, 518
 lithology, Lower Cretaceous, 579
 Lower Cretaceous, nannofossils, 221
 mineralogy, Lower Cretaceous, 579
 mollusks, Early Cretaceous, 283
 nannofossil preservation, 34
 nannofossils, Miocene, 52
 preservation, 227
 sedimentation rate, 34
 siderite, 227
 Holes 397 and 397A, correlation between, 29
 Holocene, climatic fluctuations, 727
 oceanographic and climatic variables during, 727
 Holococcoliths, 227
 defined, 53
 Holocrystalline pyroxene-plagioclase basalt fragments, constituents of volcaniclastic sands, 699
 Humic compounds, 524
 Hyalobasalt fragments, constituents of volcaniclastic sandstones, 699
 Hyaloclastites, 780
 Hydrocarbon generation, Hole 397A, 518
 migration, Black Sea, 535
 potential, 775
 source rocks, 522
 traps, 806
 Hydrocarbons, extraction of, 514
 generation of, 585
 Ice-cap stratigraphy, North America, 731
 Northern Europe, 731
 Ice-rafter material, North Atlantic, 443
 Iceland, glacial tills in, 443
 Igneous components, Clinopyroxene, volcaniclastic sandstones, 710
 feldspar, volcaniclastic sandstones, 710
 olivine, volcaniclastic sandstones, 710
 titanomagnetite, volcaniclastic sandstones, 710
 rock fragments, alkalic microgabbros, volcaniclastic sandstones, 710
 crystalline basalt, volcaniclastic sandstones, 710
 Incipient diagenesis in sediments of the lower Miocene 35
 Increased circulation along the Moroccan Continental Rise, 790
 Inertinite type organic matter, 506, 511

- Initial sedimentation phase, early to middle Miocene, 36
 Intensification of thermohaline circulation in the Atlantic Ocean, 36
 Intertropic Convergence Zone (ITCZ) in the Sahel belt, 802
 Ionian Basin, Site 125, 457
 Isomeric alkanes, 534
 Isoprenoids, 542
 Isotopic and carbonate evidence of Pliocene paleotemperatures, 402
 composition in benthic and planktonic foraminifers at Site 132, 455
 curve, 680
 - Gulf of Mexico, 735
 - Meteor* cores, 735
 - benthic foraminifers, 731
 - Niger Delta, 735
 - Panama Basin, 735
 - South Atlantic Ocean, 735
 evidence, Antarctic ice sheet, Pliocene, reduction in size during Pliocene, 443
 North Atlantic deep-water temperature variations, late Miocene, 444
 paleotemperatures, late Miocene, 453
 Isthmus of Panama, 785
 Jaramillo event, 57, 69, 472, 475, 479
 - magnetic event, 443
 Jreibichat Formation, 497, 808, 811
 Jurassic, Site 397, sedimentation rates, 808
 Jurassic/Cretaceous sedimentation, Site 397, 782
 Kaolinite, 600
 Kapitean event of New Zealand, marine regression associated with, 444
 - stage of New Zealand, late Miocene regression, 453
 Katagenesis stage, carbonification, 587
 Kerogen type, 513
 Kerogen fraction, 524
 Ketones, 541
 La Palma Island, 794
 Labrador Sea, Site 112, 66
 Lagoonal facies, 809
 Lanzarote Island, 493
 Large-scale environmental fluctuations, effects on speciation during the Pleistocene, 443
 Late Cretaceous, transgression, Aaiun Basin, 783
 - black shales, Gaada Formation, 804
 Cretaceous-Paleogene subsidence history, Site 397, 809
 Hauterivian, sedimentation rates, 80
 - sediments, 782
 Miocene, Antarctic glaciation, 36
 - dissolution event, 449
 - effect on sediment transport, Antarctic glaciation, 661
 - extent, Antarctic ice sheet, 664
 - isotopic evidence, North Atlantic deep-water temperature variations, 444
 - regression, Kapitean stage of New Zealand, 453
 - sedimentation rate, 449
 - Site 369, coarse fraction analyses, 663
 Neogene, carbonate content, Site 397, 671
 - climatic history, 433
 - environmental evolution, 447
 - grain-size, Site 397, 671
 - paleoenvironment, 391
 - Site 397, 671
 - sedimentation rate, 391, 679, 811
 - Site 397, 671
 Oligocene, regression, 83
 - Site 369, sedimentation rates, 615
 - to early Miocene, sea-level fall, 785
 Pleistocene, Arctic glaciation, 662
 - Climatic fluctuations, oxygen-isotopic record evidence, 797, 727, 733
 - correspondence to North American glacial cycles, 733
 northwestern African continental rise, sedimentation rates, 734
 oceanographic and climatic variables during, 727
 oceanography and climate, 734
 sedimentation rate, 457
 - rates, 85, 737
 Late-early to middle Miocene Canary Islands uplift event, 37
 Le Castella section, Calabria, 456
 Lebtaina Formation, 811
 Leg 1, Blake-Bahama Basin, nannofossils, 227
 Leg 42B,C1 to C5 hydrocarbon analysis, 47
 Leg 47A, core disturbance, 7
 - sediment classification, 11
 Lipid geochemistry, 541
 Lipids, Eolian origin of, 541, 543
 - Site 397, 541
 Liptinite type, organic matter, 506
 Lithology, Lower Cretaceous, Hole 397A, 579
 - Site 397, Lower Cretaceous, 560
 Lithostratigraphy, Site 397, 25
Lithraphidites bollii to *Calicalathina oblongata* zones, 564
 - Lithraphidites bollii* Zone, 66, 226
 Lower Slope anticline, 493
 - structure, 491
 Lower and middle Miocene, cyclic sediment transport, 695
 - grain morphology, sandy sediments, continental slope of northwestern Africa, 683
 - grain-size, sandy sediments, continental slope of northwestern Africa, 683
 Cretaceous, Hole 397A, lithology, 579
 - slope/upper rise region off Cape Bojador, 67
 - to middle Miocene, submarine canyons, 792
 - to Middle Cretaceous facies of Site 369, 560
 - Site 416, 560
 Lower Cretaceous biostratigraphy, Site 397, foraminifers, 257
 - depositional environment, Site 397, foraminifers, 257
 faunas, Alps, 582
 - North Caucasus, 582
 - geochemistry of clays, 583

- Hole 397A, biostratigraphy, 581
mineralogy, 579
- Lithology, Site 397, 560
nannofossils, Hole 397A, 221
northern Germany, sediments in, 566
paleoecology, Site 397, foraminifers, 257
pelagic carbonates and shales, Cape Verde Islands, 794
sedimentary structures, Site 397, 560
sedimentation rates, 487
sediments, 561
organic carbon values, 560
plant fragments, 560
siderite, 566
Site 397, diagenesis of, 567
distal delta facies accumulation, 260
silty clays, X-ray diffraction studies, 579
- Site 397, biostratigraphy, 561
slope sediments, escarpment, 491
Subsidence rates, 488
Lower Miocene shales, 523
Lower Miocene/upper Hauterivian, stratigraphic hiatus, 78, 221, 782
Lower Miocene/Hauterivian stratigraphic hiatus, 497, 742, 782, 803, 805
stratigraphic hiatus, change in $iC5/nC5$ ratio at, 534
stratigraphic hiatus, mean reflectance values bracketing, 518
stratigraphic hiatus, causes, 782
Lower Miocene/Oligocene, Site 369, stratigraphic hiatus, 783
Luminescent-microscopic analysis, organic matter, 594
Maceral types, 514
Macrofossils, 29, 32, 36, 84
ammonites, 30, 34
pelecypods, 30, 34, 59, 659
preservation of, 78
Site 397, 260, 561, 653
Magma evolution, Gran Canaria Island, 796
Magnetic anomalies *M-3* and *M-5*, 68
anomalies *M-0* to *M-25*, 66
evidence, paleocurrent directions, 481
sediment transport, 481
fabric measurements, 481
polarity, paleomagnetics, 69
reversal polarity stratigraphy in passive continental margin setting, 463
stratigraphy, sediments, 68
evidence, sedimentation rates, 471
Site 397, 470
susceptibility determinations, 464
Magnetostratigraphic evidence, biostratigraphy, 68
Mammoth and Kaena events, 456
Marine regression associated with Kapitean event of New Zealand, 444
terraces, Canary Islands, 66
on Gran Canaria Island, 85
Mass spectrometric analyses, 541
tetrapyrrole pigment content, 547
Maturation, organic matter, 805
Maturity of organic matter, Site 397, 596
- Matuyama Paleomagnetic Epoch, 57, 69, 383, 386, 391, 405, 456, 471, 479
Mauretanides, 80, 779
Mazagan Plateau, 808
Mean reflectance values bracketing stratigraphic hiatus lower Miocene/Hauterivian, 518
Mediterranean evaporite, volume of, 453
Messinian salinity crisis, 318, 449, 679
Miocene salinity crisis, 375
salinity crisis, 455
sapropelic sediments, 597
Sea, euxinification of, 451
Melt-water-related peaks, oxygen-isotope results, 728
Mesozoic/Cenozoic, Cape Bojador continental margin, subsidence history, 807
margin paleobathymetry, 807
Messinian clay sediments from Sicily, 553
Messinian salinity crisis, 85, 318, 442, 449, 679
Metadoga, new nannofossil genus, 229
mercurius, new nannofossil species, 230
Metalloporphyrins, Bahama Basin and Bermuda Rise, 550
Black Sea, 550
from the Norwegian Sea, 547
from the Norwegian-Greenland Sea, 548
odd molecular weight, 550
Meteor cores, 727
benthic foraminifers, 731
faunal fluctuations, 731
isotopic curves, 731
carbon-isotope record, 735
coarse fraction analyses, 734
foraminifers, 731
foraminifers, carbon-isotope results, 728
Oxygen-isotope results, 728
isotopic curves, 735
oxygen-isotope stratigraphy, 734
sedimentary record, 734
Methane, 615, 806
biogenic, 531, 532
biological, 804
carbon-13 values, 533
Site 397, 804
Microfaulting, sediment structures, 781
Microstaurus conus, new nannofossil combination, 230
Mid-Cretaceous organic-rich shales, Aaiun-Tarfaya Basin, 804
Site 369, 805
Mid-Miocene uplift of Canary Island volcanoes, 792
Middle to early Miocene, turbidite sandstone beds, Site 397, 806
lower Miocene, turbidites, 86
Middle Cretaceous subsidence of offshore depositional sites, 261
transgression, 261, 811
Middle Miocene, Canary Island volcanism, 86, 87
environmental crisis, 64
Fuerteventura volcanic debris flows, seismic reflection profiles, 795
petrochemistry of volcaniclastic sandstones, 699
petrography of volcaniclastic sandstones, 699

- sedimentary cycles, 693
 sedimentation rate, 764
 rates, 84, 613
 stratigraphic hiatus, Zones NN 7 to NN 9, 296
 to Quaternary, Sites 369 and 397, volcanism, 796
 upwelling with buildup of eastern Antarctic ice cap, 798
 volcanic apron of, Gran Canaria Island, 497
 Middle/upper Eocene, Stratigraphic hiatus Site 369, 783
 Migmatic texture of mudstones, 29, 780
 Miliolids, Site 397, 260
 Mineral carbon content, Site 397, 523
 Mineralogy, Lower Cretaceous, Hole 397A, 579
 Site 141, sediments, 644
 Miocene, glacial expansion in Antarctica, 455
 Canary Islands volcanism, 615
 earthquake activity, Atlas Mountains, 615
 foraminiferal stratigraphy, Site 397, 317
 Site 366, 341
 foraminifers, 59, 317
 Hole 397A, nannofossils, 52
 mollusks, 63
 nannofossils, 52, 291
 oxygen-minimum zone, North Atlantic, 520
 salinity crisis, Mediterranean, 375
 stratigraphy, Site 372, foraminifers, 321
 Miocene/Pliocene boundary, 443
 boron geochemistry, 553
 foraminifers, 59, 318, 327, 341, 673
 nannofossils, 386
 Site 397, 386
 X-ray diffraction studies of sediments, 553
 Mississauga Formation, prodelta mudstones, 567, 810
 Mississippi delta, prodelta sediments, 564
 modern prodelta sediments of, 260
 Modern near-shore and on-shore sedimentary environments, northwestern Africa, 736
 prodelta sediments of the Mississippi Delta, 260
 Mollusks, Cretaceous, 64
 Early Cretaceous, Hole 397A, 283
 Miocene, 63
 systematics, 284
 Montmorillonite, 752
 Mordenite, zeolites, 747
 Moroccan coastal basins, Aptian/Albian, 811
 Continental Rise, increased circulation along, 790
 Morocco Basin, 804
 Mudstones, migmatic texture of, 29, 780
 Multichannel seismic profiles, 19, 87, 493
 N-alkane distribution, 508
 N-alkanes, 515, 526, 541
 trends, 585
 N-fatty acids, 541
 N-methyl ketones, Black Sea, 542
 Nannofoossil preservation, Hole 397A, 34
 Nannofoossils
 biostratigraphy, 67
 Cenozoic, abundance, 296
 preservation, 296
 Cretaceous, 52
 Diloma, new genus, 228
 placinum, new species, 228
 primitiva, new combination, 228
 Hole 397A, Lower Cretaceous, 221
 Holococcoliths, defined, 53
 Leg 1, Blake-Bahama Basin, 227
 Metadoga, new genus, 229
 mercurius, new species, 230
 Microstaurus conus, new combination, 230
 Miocene, 52, 291
 Hole 397A, 52
 Miocene/Pliocene boundary, 386
 Neogene sedimentation rates based on, 64
 and Quaternary, 289
 NN 18/NN 19 boundary, 51
 NN 19/NN 20 zonal boundary, 472
 Octopodorhabdus plethotretus, new species, 230
 Parhabdolithus swinnertonii, new combination, 231
 Pleistocene, 51
 Pliocene, 51
 preservation, Hole 397A, 227
 Quaternary, 295
 reworked, 298
 Rhabdolekiskus parallelus, new species, 232
 Site 397, 51
 systematics, 228
 Trapezopentus, new genus, 232
 sarmatus, new species, 232
 Watznaueria fasciata, new species, 232
 Zonation, Anaurolithus primus Zone, 380
 Calcicalathina oblongata zone, 66, 226
 Ceratolithus acutus Zone, 380
 rugosus Zone, 380
 tricorniculatus and Ceratolithus rugosus zones (NN 12-13), 293
 Zone (NN 12) to Discoaster brouweri (NN 18), 64
 Cretarhabdus loriei Zone, 226
 Cruciellipsis cuvillieri Zone, 53
 Discoaster asymmetricus Zone (NN 14), 294
 berggrenii zone, 379
 Subzone, 380
 brouweri Zone, (NN 18), 51, 294, 381
 calcaris Zone (NN 10), 52, 293
 druggii Zone (NN 2), 64, 221, 289, 291
 exilis Zone (NN 6), 52, 293
 hamatus Zone, (NN 9) 293, 377
 to Discoaster quinqueramus Zone (NN 11)
 64, 298
 kugleri Zone (NN 7), 52, 293
 neohamatus Zone, 377
 pentaradiatus Zone (NN 17), 51, 294, 381
 quinqueramus Zone (NN 11), 293
 surculus Zone, (NN 16), 51, 294, 381
 tamalis Zone, 384
 Emiliania huxleyi Zone (NN 21), 51, 289, 296
 Gephyrocapsa oceanica Zone (NN 20), 51, 289, 296
 Helicopontosphaera ampliaperta Zone (NN 4), 52, 293, 296, 298

- sellii* Zone, 295
Lithraphidites bollii to *Calcicalathina oblongata*
 zones, 564
Lithraphidites bollii Zone, 66, 226
Pseudoemiliania lacunosa Zone (NN 19), 51, 295
 to *Emiliania huxleyi* zone (NN 21), 64
Reticulofenestra pseudoumbilica Zone (NN 15),
 51, 294, 381, 386
pseudoumbilica/Discoaster surculus zonal
 boundary, 382
Sphenolithus belemnos Zone (NN 3), 52, 291, 296
 to *Helicopontosphaera ampliaperta* Zone
 (NN 4), 64
distantus Zone (NP 24), 67
heteromorphus Zone (NN 5), 52, 293
 to *Discoaster kugleri* Zone (NN 7), 64, 296
Triquetrorhabdulus carinatus Zone (NN 1), 291
rugosus Zone, 380
Tubodiscus jurapelicus Zone, 222
 Zone NN 11, 52, 380
 NN 13, 341
 NN 14, 64
 NN 16, 381
 NN 17, 381
 NN 2, 67, 782, 790
 NN 9, 52, 63
 NP 24, 782
 Zones NN 11 to NN 14, 62
 NN 13 and NN 12, 52
 NN 15 to NN 19, 62
 NN 2, 63
 NN 3 to NN 7, 63
 NN 7 to NN 9, Stratigraphic hiatus, Middle Miocene, 296
 NN 9 and NN 10, 611
 NN 8, 52
 NN 8 to NN 11, 60
 Natural remanent magnetization (NRM), 86
 Sediments, 68
 Site 397, 464
 Variation downhole, 475
 Neocomites, ammonites, 283
 Neogene, autochthonous sequence, Site 397, sediments,
 647
 and Quaternary benthic foraminifers, 60
 nannofossils, 289
 climates, reconstructed from clay fraction analysis,
 643
 reconstructed from coarse fraction analysis, 643
 climatic conditions, reconstructed from coarse fraction analysis, 662
 evolution, Canary Island volcanism, 703
 Paleobathymetry, Site 397, 809
 sedimentation rates, 296
 based on foraminifers, 64
 nannofossils 64
 sediments, 774
 sediments in region of Cape Bojador, 298
 sequence, Site 369, 647
 Site 397, sedimentation rate, 520
 rates, 64, 647
 Subsidence, Site 397, 809
 volcanic uplift of Canary Archipelago, 713
 Neopentane, 534, 537
 Nepheloid layer, particulate matter suspended in, 504
 layers, 564
 New nannofossil combination, *Diloma primitiva*, 228
 Microstaurus conus, 230
 Parhabdolithus swinnertonii, 231
 New nannofossil genus, *Diloma*, 228
 Metadoga, 229
 Trapezopentus, 232
 New radiometric age determinations of Pliocene/Pleistocene boundary, 457
 New foraminiferal species, *Favusella stiftia*, 276
 New nannofossil species, *Diloma placinum*, 228
 Metadoga mercurius, 230
 Octopodorhabdus plethotretus, 230
 Rhabdoekiskus parallelus, 232
 Trapezopentus sarmatus, 232
 Watznaueria fasciata, 232
 Newfoundland continental margin, Early Cretaceous, 567
 Newfoundland shelf basin, 810
 Nickel porphyrins, 548
 Niger delta, 804
 isotopic curves, 735
 prodelta sediments, 561
 North and South Atlantic Ocean, stratigraphic hiatus,
 Oligocene, 782
 North African continental margin, sedimentation rate, 448
 North America, ice-cap stratigraphy, 731
 North Atlantic, deep-water, 665
 oxygen content, 422
 Pliocene, oxygen isotopic composition, 443
 temperature variations, late Miocene, isotopic evidence, 444
 ice rafted material, 443
 Miocene, oxygen-minimum zone, 520
 North Caucasus, Lower Cretaceous faunas, 582
 Northern guests, criterion of, ostracodes, 457
 Northern Europe, ice-cap stratigraphy, 731
 Northern Germany, sediments in Lower Cretaceous, 566
 Northern Hemisphere ice-sheet development, glacio-eustatic regression, 85
 Northern Italy, Po Plain, 398
 Northwest African continental margin off Cape Bojador, 617
 margin, debris flows from, 35
 Site 397, 17
 passive margin, 603, 616
 Northwestern Africa, geologic history of Cape Bojador continental margin, 773
 modern near-shore and onshore sedimentary environments, 736
 African continental rise, sedimentation rates, late Pleistocene, 734
 continental margin, geologic history of, 775

- margin, submarine slides and slumps, Quaternary, 792
 Norwegian Sea, metalloporphyrins from, 547
 Norwegian-Greenland Sea, metalloporphyrins from, 548
 Nova Scotian continental margin, Early Cretaceous, 567
 shelf basin, 810
 Nunivak and Cochiti events, 384
 Oceanic circulation, Hautervian, 568
 Oceanographic and climatic variables during the Holocene, 727
 the late Pleistocene, 727
 Oceanography and climate, late Pleistocene, 734
Octopodorhabdus plethotretus, new nannofossil species, 230
 Odd molecular metalloporphyrin series, 551
 weight, metalloporphyrins, 550
 Odd/even carbon preference, organic matter, 516, 542
 Offshore depositional sites, Middle Cretaceous subsidence of, 261
 wells, thermal gradient, 70
 Olduvai event, 69, 456, 471, 479
 Oligocene
 climatic deterioration, 788
 sea-level regression, 785
 sedimentation rates, 785
 North and South Atlantic Ocean, stratigraphic hiatus, 782
 Olivine, volcaniclastic sandstones, igneous components, 710
 Onset of glaciation, Antarctic ice cap, 785
 Ooze/chalk transition at Site 167, 448
 Opal-indicated upwelling maxima, 735
 Opal-A, 742
 Opal-CT, 87, 742, 745, 746, 747, 780, 797
 Open basin Tethyan environment, 284
 Operations, Site 397, 21
 Optical emission spectrography analyses, boron, concentration in sediments, 553
Orbulina suturalis/Globorotalia peripheroronda Zone (N.9), 59
 Organic and inorganic particles, settling rates for, 505
 carbon, value, 665
 analyses, Site 397, 47
 content, Site 397, 524
 variability in determinations, 506
 facies, 509
 geochemistry, C1 to C5 hydrocarbon analysis, 47
 matter, algal type, 506, 511
 and hydrocarbon gases, diagenesis, 86
 biodegradation of, 503
 diagenesis of, 775
 even/odd carbon preference, 542
 fusinite type, 514
 inertinite type, 506, 511
 in deep ocean sediments, transport and depositional mechanisms, 503
 in the world's oceans, cycling of, 503
 liptinite type, 506
 luminescent-microscopic analysis, 594
 marine derived, 508
 maturation, 805
 odd/even carbon preference, 516, 542
 Site 397, maturity of, 596
 terrestrial derivation, 508, 518
 types of, 506
 vitrinite type, 506, 511, 514
 maturity, sediments, Site 397, 505
 Time-temperature effects, 506
 Vitrinite reflectance, 518, 522
 petrography, Site 397, 49
 productivity at Site 62.1, 448
 Organic-rich shales, pyrolysis assay of, 524
 Site 397, 523
 Miocene sediments from the Vørning Plateau, 509
 Ostracodes, 455, 582
 Site 397, 260, 653
 Ostracodes, Northern guests, criterion of, 457
 Oued Draa River, 643
 Oxidizing-reducing conditions, 587
 Oxygen and carbon isotope stratigraphy, benthic foraminifers, 433
 content, North Atlantic deep-water, 422
 depletion, effect on litho- and biofacies, 565
 isotopic composition, North Atlantic deep-water, Pliocene, 443
 composition changes, Site 284, 443
 Oxygen-isotope curves, continental margin cores, 728
 data, climatic cycles identified with, 727
 record effects, upwelling, 735
 results, melt-water-related peaks, 728
 Meteor cores, foraminifers, 728
 stratigraphy, *Meteor* cores, 734
 Oxygen-isotopic record evidence, late Pleistocene, Climatic fluctuations, 797
 Oxygen-isotope curve, surface-water, 735
 Oxygen-minimum layers, 615
 zone, North Atlantic, Miocene, 520
 Oxygen-18 isotope curve, 419
 values, Site 397, 85, 393, 664
 Paleobathymetric estimations, Site 397, 764
 Paleobathymetry, benthic foraminifers, 62
 Mesozoic/Cenozoic, Cape Bojador continental margin, 807
 Site 397, Neogene, 809
 Paleoclimates, Eocene through Oligocene, 785
 Sahara Desert, 802
 Paleoclimatic record, Site 397, 391
 Paleocurrent directions, magnetic evidence, 481
 Paleodepth, Canary Island channel, 795
 Paleodepths, Site 397, Early Cretaceous, 809
 Paleoecology, Site 397, foraminifers, Lower Cretaceous 257
 Paleoenvironment, Early Cretaceous, 561
 evolution, Site 369, 77
 late Neogene, 391
 sedimentary history, Cape Bojador continental margin, Early Cretaceous, 559

- Site 397, 782
 late Neogene, 671
 Paleogene sedimentary rocks, fibrous clay source, 643
 Paleomagnetic Epoch 11, 69
 5, 341, 384, 451, 679
 6, 386, 479
 7, 59, 69, 384
 Epochs 16 through 19, 69
 5 and 6, 444
 9/10, 69
 measurements, chemical overprinting, Sites 379 and 380, 464
 reversal record, Site 397, 479
 studies, Site 397, 463
 Paleomagnetics, magnetic polarity, 69
 Site 397, 68
 volume susceptibility, 69
 Paleopressures, sedimentary basin, 761, 768
 Paleotemperature curves based on foraminifers, 735
 index 2, 733
 Paleotemperatures, late Miocene, isotopic evidence, 453
 Pliocene, isotopic and carbonate evidence, 402
 sedimentary basin, 761, 768
 Paleowinds, Trade Winds in the Wheel-Round latitudes, 737
 Panama, Isthmus of, 785
 basin, isotopic curves, 735
 Paratethys, euryhaline fauna from, 455
Parhabdolithus swinnertonii, new nannofossil combination, 231
 Particle-by-particle displacement, sediment transport, 737
 Particulate matter suspended in nepheloid layer, 504
 organic matter concentrations in waters of the Caribaco Trench, 506
 Passive continental margin, in the eastern North Atlantic during the Mesozoic, geologic history, 257
 of northern Africa, Quaternary, 614
 off northwest Africa, stratigraphy and evolution of, 741
 setting, magnetic reversal polarity stratigraphy in, 463
 subsidence mechanisms, 810
 margins, stratigraphy and evolution of, 19
 evolution of, 617
 Pelagic carbonate sediments, diagenesis, 448
 nannofossil marls, Site 369, Upper Cretaceous, 809
 Pelecypods, macrofossils, 30, 34, 59, 659
 Periodicity of sedimentogenesis, 692
 Permeability, physical properties, sediments, 761
 Petrochemistry of volcaniclastic sandstones, middle Miocene, 699
 Petrography of volcaniclastic sandstone, 707
 sandstones, middle Miocene, 699
 Petroleum formation, C2 to C4 hydrocarbons diagnostic of, 531
 generation, sediments, 86
 potential, African margin, 78
 Phillipsite, 746, 748, 751
 Phosphate grains, Site 397, coarse fraction analysis, 660
 Phosphorite occurrences, upwelling, 664
 Phyllosilicates, authigenic, 748
 Sites 397 and 398, 741
 Physical and thermodynamic properties of sediments, temperature effects on, 766
 Physical properties, 13
 fluids, density, 761
 viscosity, 761
 sediments, compaction, 761
 permeability, 761
 porosity, 761
 Site 397, 37
 compared with Site 397 GRAPE measurements, 766
 Physiographic setting, Site 397, 19
 Phytane, 516, 520
 Plant fragments, Lower Cretaceous, 560
 Site 397, 560
 Pleistocene, carbon isotope fluctuations, 443
 ice conditions, Arctic Ocean, diatom evidence, 479
 large-scale environmental fluctuations, effects on speciation, 443
 nannofossils, 51
 sedimentation rate, 442
 Pliocene, foraminifers, 58
 isotopic and carbonate evidence, paleotemperatures, 402
 nannofossils, 51
 oxygen isotopic composition, North Atlantic deep-water, 443
 reduction in size during Pliocene, isotopic evidence Antarctic ice sheet, 443
 Pliocene/Pleistocene boundary, 443, 456
 foraminifers, 57
 new radiometric age determinations of, 457
 stratotype at Calabria, 456
 Pliocene/Quaternary sedimentation rate, 764
 Poe Plain, Northern Italy, 398
 Porcelaneous mudstones, early Miocene, 751
 Pore pressure, computer modeling, Site 397, 765
 water analyses, Site 397, 49
 Porosity, physical properties, sediments, 761
 Possible causes of continental margin destruction, 785
 Potassium-argon age determinations of volcanic samples from the Canary Islands, 666
 Pre-early Miocene sculpturing event, 611
 Pre-late Oligocene or early Miocene sculpturing event, 36
 Precambrian Anti-Atlas Uplift-South Atlas Fault, 779
 Requibat High, 19
 Present influence of upwelling, Site 397, 665
 Trade Wind belt off Cape Blanc, 665
 Pristane, 516, 520
 to phytane ratio, 542
 Prodelta, 596
 mud environment, Site 397, 809
 muds, 794
 mudstones, Mississauga Formation, 810
 sediments, Guadalupe delta, 564
 Mississippi delta, 564
 Niger delta, 561
 setting, 509

- Pseudoemiliania lacunosa* Zone (NN 19), 51, 295
 to *Emiliania huxleyi* Zone (NN 21), 69
- Puerto Rico Trench, Recent, 84
- Pull-apart structures, sediment structures, 781
- Pyrite, 781
 Site 397, 648
 coarse fraction analyses, 658
 nodules, Site 397, 553
 noise, effect on grain-size data validity, 677
- Pyrolysis, 505
 assay of organic-rich shales, 524
- Pyrolysis-fluorescence techniques, Site 397, 48
- Quartz, volcaniclastic sandstones, 712
 grains, tracers of arid climates, 641
- Quartz, volcaniclastic sandstones, 712
 grains, tracers of arid climates, 641
- Quartz-cherts, 751
- Quaternary changes in Atlantic deep-water circulation, 419
 climate, Senegal, 662
 climates, reconstructed from coarse fraction analyses, 661
 foraminifers, 57
 nannofossils, 295
 northwestern African margin, submarine slides and slumps, 792
 passive continental margin of northern Africa, slumps and debris flows, 614
 Site 397, calcium carbonate dissolution, 665
- Quaternary/Pliocene boundary, coarse fraction analysis, 660
- Radiolarian, Site 397, 653
- Recent, Puerto Rico Trench, 84
 Niger prodelta, 80
- Recycled organic matter, spores, 514
- Reddish quartz, Senegal Delta, 736
- Reflectivity, vitrinite-huminite, 511
- Refraction velocities, 491
- Refractive indexes of siliceous organisms, 742
- Regression, Kapitean stage of New Zealand, late Miocene, 453
 late Oligocene, 83
- Relict particles, coarse fraction analyses, Site 397, 658
- Requibat Massif, 80, 567, 779, 807
 Uplift, 77
- Reservoir Beds, potential, Site 397, 806
- Reticulofenestra pseudoumbilica* Zone (NN 15), 51, 294, 381, 386
pseudoumbilica/Discoaster surculus zonal boundary, 382
- Reunion event, 457
- Rhabdolekiskus parallelus*, new nannofossil species, 232
- Rhyolitic ash layers, Sites 369 and 397, 811
 eruption on Gran Canaria Island, 85
- Rip-up clasts, sediment structures, 781
- Roll-up structures, sediment structures, 781
- Sahara coastal margin, climatic history of, 737
 Desert, 661
 climatic fluctuations, carbon-14 record, 736
 paleoclimates, 802
- Sahel belt, intertropic Convergence Zone (ITCZ) in the, 802
- lateritic belt, Sahara Desert, dust-derived sediments, 803
 zone, 737
- Salinity fluctuations, 586
- Samlat Formation, 20, 811
- Santerno section of northern Italy, 457
- Schlanger and Douglas diagenetic model, 448
- Scotian Shelf, 810
 Subsidence rates, 809
 foraminiferal genus *Favusella*, 273
- Sculpturing event, pre-early Miocene, 611
 pre-late Oligocene or early Miocene, p. 36
- Sea-level fall, late Oligocene to early Miocene, 785
 regression, Oligocene, 785
- Seaward transport of terrestrial organic matter, 503
- Sediment classification, Leg 47A, 11
 diagenesis, electron microscopy studies, 742
 X-ray diffraction studies, 742
- load, Senegal River, 736
 section, Site 369, 20
 structures, decollement zones, 781
 folds, 781
 microfaulting, 781
 pull-apart structures, 781
 rip-up clasts, 781
 roll-up structures, 781
 transport, 85, 617, 644, 660, 692
 by slumping, 509
 turbidity currents, 509
 downslope displacement, 737
 lower and middle Miocene, cyclic, 695
 magnetic evidence, 481
 particle-by-particle displacement, 737
- Sedimentary rock fragments, volcaniclastic sandstones, 710
 basin, compaction, 761
 depositional environment, Early Cretaceous, 585
 paleopressures, 761, 768
 paleotemperatures, 761, 768
 subsidence, 761
 unstable salinity regime, depositional environment, Early Cretaceous, 587
 cycles, middle Miocene, 693
 history, Cape Bojador continental margin, Early Cretaceous, 559
 mélange deposits, 35
 processes, downslope displacement, 55
 reworking from older formations, 55
 Site 397, 782
 record, *Meteor* cores, 734
 sequence, Aaiun Basin, 559
 structures, 26, 27, 30
 fold structures, 612
 Site 397, Lower Cretaceous, 560
 volcaniclastic sandstones, 28
- Sedimentation of ferriferous carbonate bands, 586
 rate, Albian to Upper Cretaceous, 488
 early Miocene, 764
 Hauterivian, Site 397, 564
 Hole 397A, 34
 late Miocene, 449
 late Neogene, 391, 679

- late Pleistocene, 457
 middle Miocene, 764
 Neogene, Site 397, 520
 North African continental margin, 448
 Pleistocene, 442
 Pliocene/Quaternary, 764
 Site 397, late Neogene, 671
 Tertiary, Site 397, 518
 upper Miocene, 764
 rates based on foraminifers, Neogene, 64
 nannofossils, Neogene, 64
 Cenomanian to Turonian, 568
 Cretaceous, 65
 Early Cretaceous, 782
 paleomagnetic evidence, 475
 Miocene, 612
 Jurassic, Site 397, 808
 late Hauterivian, 80
 Neogene, 811
 Oligocene, Site 369, 615
 Pleistocene, 85, 737
 northwestern African continental rise, 734
 Lower Cretaceous, 487
 magnetic stratigraphy evidence, 471
 middle Miocene, 84, 613
 Neogene, 296
 Site 397, 64, 647
 Oligocene, 785
 Upper Neogene, 751
 Sedimentogenesis, periodicity of, 692
 Sediments, Baie du Levrier, 736
 chemical analyses, Site 397, 49
 compaction, physical properties, 761
 Cretaceous (post Hauterivian), Site 369, 782
 late Hauterivian, 782
 Lower Cretaceous, 561
 age of, 226
 magnetic stratigraphy, 68
 mineralogy, Site 141, 644
 Miocene/Pliocene boundary X-ray diffraction studies of, 553
 natural remanent magnetization (NRM), 68
 Neogene, 774
 autochthonous sequence, Site 397, 647
 organic carbon values, Lower Cretaceous, 560
 permeability, physical properties, 761
 petroleum generation, 86
 plant fragments, Lower Cretaceous, 560
 porosity, physical properties, 761
 shallow water particles, Site 369, 661
 siderite, Lower Cretaceous, 566
 Site 369, 487
 397, diagenesis of, 567
 organic maturity, 505
 terrigenous components, Site 397, 641
 thermal properties, 761
 Seismic interpretation, Cape Bojador slope, 491
 reflection profiles, Fuerteventura volcanic debris flow, Middle Miocene, 795
 correlated with drilling results, 70
 of West Saharan continental margin between Cape Bojador and Cape Blanc, 20
 reflectors, 71
 stratigraphic correlation of, 73, 87
 sequences, Site 397, 485
 Selection of Site 397, 20
 Senegal, Quaternary climate, 662
 Basin, 804
 Delta, Reddish quartz in, 736
 River, 802
 sediment load, 736
 Sepiolite, 600
 Sequence of events in Neogene history of, African margin off Cape Bojador, 36
 Sequia el Hamra dry river, 497
 Serpentine, 713
 Serpulidae, 660
 Site 397, 653
 Settling rates for organic and inorganic particles, 505
 Shallow water particles, Site 369 sediments, 661
 sediment supply, Site 397, 659
 Shallow-water sediment sequences, Tarfaya Basin, 794
 Shear strength, Site 397, 45
 Shield building, Canary Island, basalt, 714
 Sicily, Messinian clay sediments from, 553
 Siderite, 34, 581
 Hole 397A, 227
 Lower Cretaceous, sediments, 566
 layers, 603, 774
 Boudinage-structure, 560
 nodules, 561
 in near-shore Holocene muds, Strait of Malakka, 566
 Sideromelane shards, ash layers, 711
 Silica, authigenic, 751
 cementation by, 747
 diagenesis, 87, 750, 751
 Sites 397 and 398, 741
 Siliceous organisms, diagenesis of, 742
 preservation of, 742
 refractive indexes of, 742
 transformation to opal-CT quartz, 742
 Silty clays, X-ray diffraction studies, Lower Cretaceous, 579
 Site 105, foraminiferal, genus, *Favusella*, 274
 112, Labrador Sea, 66
 125, Ionian Basin, 457
 132, Isotopic composition in benthic and planktonic foraminifers at, 455
 Tyrrhenian Basin, 457
 138, Contourites, 793
 West of the Canary Islands, 790
 139, Eolian sand turbidites, 793
 141, Sediments, mineralogy, 644
 144, Demerara Rise, 566
 foraminiferal genus, *Favusella*, 274
 147, Cariaco Trench, 531
 167, Equatorial Pacific Ocean, chalk diagenesis, 674
 ooze/chalk transition at, 448
 284, oxygen isotopic composition changes, 443

- 366, Miocene foraminiferal stratigraphy, 341
 367, Benthic foraminifers, 60
 Cape Verde Basin, 221
 368, Cape Verde Rise, contourites, 793
 369, benthic foraminifers, 60
 calcium carbonate dissolution, upper Miocene, 665
 coarse fraction analyses, late Miocene, 663
 Lower to Middle Cretaceous facies of, 560
 Mid-Cretaceous organic-rich shales, 805
 Neogene sequence, 647
 paleoenvironment evolution, 77
 sediment section, 20
 sedimentation rates, late Oligocene, 615
 sediments, 487
 Cretaceous (post Hauterivian), 782
 shallow water particles, 661
 Stratigraphic hiatus, Lower Miocene/Oligocene, 783
 middle/upper Eocene, 783
 submarine canyon closest to, 792
 Upper Cretaceous pelagic nannofossil marls, 809
 and Paleogene sediments, 221
 volcanic material, eolian transport of, 666
 370, Cape Verde Basin, 221
 372, foraminifers, Miocene stratigraphy, 321
 Site 397, acoustic horizons, 448
 acoustic velocity, 41
 Agadir-Essaouira Region, correlated with offshore prodelta facies, 261
 ash layers, 703
 background and objectives, 19
 benthic foraminifers, 419
 biostratigraphy, 51
 and magnetostratigraphy, 85
 Lower Cretaceous, 561
 bulk physical properties, 44
 properties of sediments, 45
 calcium carbonate dissolution, Quaternary, 665
 upper Miocene, 665
 Canary Islands volcanic center, effect on geothermal gradients, 508
 carbon-13 values, 86, 398, 508
 carbonate values, upper Miocene, 672
 chalk diagenesis, 674
 chlorophyll diagenesis, 547
 clay fraction analysis, 643
 mineralogy, 595
 minerals, x-ray diffraction studies of, 595
 coarse fraction analyses, glauconite, 658
 pyrite, 658
 relic particles, 658
 terrigenous material, 658
 volcanic material, 658
 analysis, 641, 647
 phosphate grains, 660
 results, summary of, 30
 composition of ash layers, 707
 computer modeling, 765
 continental climate and deep-sea record, 802
 C1 to C7 hydrocarbons, 531
- depositional environments and processes, 33
 diagenesis of Lower Cretaceous, sediments, 567
 diatoms, 653
 dissolution at depth, foraminifers, 56
 distal delta facies accumulation, Lower Cretaceous sediments, 260
 downhole temperature measurements, 69
 Early Cretaceous, paleodepths, 809
 echinoderms, 653
 eolian material, 695
 sand turbidites, 644
 fish debris, 560, 653
 foraminifers, 53
 Lower Cretaceous, biostratigraphy, 257
 depositional environment, 257
 paleoecology, 257
 Fuerteventura Island, volcanic components source, 714
 geochemistry, 46
 geothermal gradient, 508, 512, 764
 Gran Canaria Island, volcanic components source, 714
 GRAPE mesurements, physical properties compared with, computer modeling, 756
 Jurassic/Cretaceous sedimentation, 782
 Late Cretaceous-Paleogene subsidence history, 809
 Neogene, carbonate content, 671
 grain-size, 671
 paleoenvironment, 671
 sedimentation rate, 671
 lipids, 541
 lithostratigraphy, 25
 Lower Cretaceous, Lithology, 560
 Sedimentary structures, 560
 macrofossils, 260, 561, 653
 magnetic stratigraphy, 470
 maturity of organic matter, 596
 methane, carbon-13 values, 804
 middle to early Miocene turbidite sandstone beds, 806
 miliolids, 260
 mineral carbon, content, 523
 miocene foraminiferal stratigraphy, 317
 Miocene/Pliocene boundary, 386
 nannofossils, 51
 natural remanent magnetization (NRM) measurements, 464
 Neogene, paleobathymetry, 809
 subsidence, 809
 northwest African margin, 17
 operations, 21
 organic carbon, content, 524
 analyses, 47
 maturity, sediments, 505
 petrography, 49
 organic-rich shales, 523
 ostracodes, 260, 653
 oxygen-18 values, 85, 393, 664
 paleobathymetry estimates, 764
 paleoclimatic record, 391
 paleoenvironment, 782

- paleomagnetic reversal record, 479
 studies, 463
- paleomagnetics, 68
- physical properties, 37
- physiographic setting, 19
- plant fragments, 560
- pore, pressure, computer modeling, 765
 water analyses, 49
- present influence, upwelling, 665
- preservation, foraminifers, 57
- prodelta mud environment of, 809
- pyrite, 648
 nodules, 553
- pyrolysis-fluorescence techniques, 48
- radiolarians, 653
- reservoir beds, potential, 806
- sedimentation processes, 782
- sedimentation rate, Hauterivian, 564
 Neogene, 520
 Tertiary, 518
- rates, Jurassic, 808
 Neogene, 64, 647
- sediments, chemical analyses, 49
 Neogene, autochthonous sequence, 647
 terrigenous components, 641
- seismic sequences, 485
- selection of, 20
- Serpulidae, 653
- shallow-water sediment supply, 659
- shear strength, 45
- smectite, 599
- stratigraphic hiatuses, 66
- summary and conclusions, 77
- Tarfaya-Aauin Region, correlated with offshore prodelta facies, 260
- thermodynamic parameters, computer modeling, 765
- Tortonian/Messinian boundary, 386
- upper Quaternary influence, upwelling, 665
- variation downhole, natural remanent magnetization (NRM), 475
- volcanic debris, clastic origin of, 714
 material, submarine transport of, 666
- volcaniclastic sandstones, 703
- Zanclean/Pizcenzian boundary, 387
- Site 398, Lower Cretaceous sediments, 596
 416, Lower to Middle Cretaceous facies of, 560
 62.1, organic productivity at, 448
 97, foraminiferal genus, *Favusella*, 274
- Sites 379 and 380, paleomagnetic measurements, chemical overprinting, 464
- 360 and 361, South Atlantic Basin, 531
 369, drilling information, 491
 369 and 397, rhyolitic ash layers, 811
 volcanism, middle Miocene to Quaternary, 796
- 379-381, Black Sea, 531
 397 and 398, phyllosilicates, 741
 silica, 741
 zeolites, 741
- Slope anticline, 19, 83, 487, 309, 810
 lower, 491, 493
- Slope-rise destruction, deep-current erosion causing, 34
- Slump deposit, 614
 structures, 34, 561, 579, 779
- Slumped zones, 781
- Slumping, 55, 64, 84, 318, 381, 560, 606, 611, 612, 615, 765, 785
 continental along near Cape Bojador, 19
 displacement of Samlat Formation rocks, 20
 episodes, 28
 sediment transport by, 509
 zones, 27
- Slumps, 782
 debris flows, Quaternary, 614
- Smectite, 84, 643, 713, 780
 Atlas Mountains source, 616
 cementation by, 747
 Site 397, 599
- Solution index, foraminifers, 671
- South Atlantic Basin, Sites 360 and 361, 531
 Ocean, isotopic curves, 735
- South Canary Basin, 615
 Island Channel, 611, 707
- Sphaeroidinellopsis acme* Zone, 455
seminulina paenedehiscense Zone, 59, 339
subdehiscens partial range Zone, 58
subdehiscens/Globigerina druryi Zone (N.13), 59
- Sphenolithus belemnos* Zone (NN 3), 52, 291, 296
 to *Helicopontosphaera ampliaperta* Zone
 (NN 4), 64
- distentus* Zone (NP 24), 67
heteromorphus Zone (NN 5), 52, 293
 to *Discoaster kugleri* Zone (NN 7), 64, 296
- Spores, recycled organic matter, 514
- Stable isotopic ratios correlated with carbonate curves, 671
- Stagnant basin conditions, 586
- Stained-quartz fractions, source of, 31
- Sterenes, 526
- Steroid hydrocarbons, 516, 521, 542
- Strait of Malakka, siderite nodules in near-shore Holocene muds, 566
- Stratigraphic correlation of seismic reflectors, 73, 87
 hiatus, Early Cretaceous, 60
 Eocene/Paleocene, 485
 seismic reflection profile evidence, 488
 lower Miocene/upper Hauterivian, 78, 221
 lower Miocene/Hauterivian, 742, 782, 803, 805
 causes, 782
 change in iC5/nC5 ratio at, 534
 mean reflectance values bracketing, 518
 lower Miocene/Oligocene, Site 369, 783
 middle Miocene, nannofossil, zones NN 7 to NN 9, 296
 middle/upper Eocene, Site 369, 783
 Oligocene, North and South Atlantic Ocean, 782
 hiatuses, Site 397, 66
- Stratigraphy and evolution of the passive continental margin off northwest Africa, 741
 evolution of passive continental margins, 19
- Subaerial deltaic muds, Aaiun/Tarfaya Basin, 78

- volcanic stage of the Canary Island Archipelago, 811
 Submarine canyon closest to Site 369, 792
 canyons, 611, 615
 lower to middle Miocene, 792
 erosion, biostratigraphic evidence for, 57
 shield of Gran Canaria Island, 796
 slides and slumps, Quaternary, northwestern African margin, 792
 transport of Site 397 volcanic material, 666
 Subsidence, Cape Bojador continental margin, 809
 history, Mesozoic/Cenozoic, Cape Bojador continental margin, 807
 mechanisms, passive continental margin, 810
 of Cape Bojador continental margin, 568
 rates, Baltimore Canyon Trough, 809
 Early Cretaceous, 809
 Lower Cretaceous, 488
 Scotian Shelf, 809
 sedimentary basin, 761
 Site 397, Neogene, 809
 Summary and conclusions, Site 397, 77
 Surface-water, oxygen-isotope curve, 735
 regime changes correlated with Atlantic deep-water circulation changes, 735
 Systematics, mollusks, 284
 nannofossils, 228
 Tachylite, volcaniclastic rock fragments, 710
 Tarfaya Basin, 283, 284
 shallow-water sediment sequences, 794
 Tarfaya-Aauin Region, correlated with offshore pro-delta facies, Site 397, 260
 Tectonics, Atlas Mountains, 663
 Temperature effects on physical and thermodynamic properties of sediments, 766
 Tenerife Island, 794
 volcanic eruption on, 85
 Terpenes, 537
 Terrestrial biomass, effect on carbon-13 values, 442
 derivation, organic matter, 508, 518
 organic matter, seaward transport of, 503
 Terrigenous input to continental margin of the eastern North Atlantic, 736
 components, Site 397, sediments, 641
 material, Site 397, coarse fraction analyses, 658
 Tertiary, Site 397, sedimentation rate, 518
 Tethyan environment, open basin, 284
 Tethys, 810
 Tetrapyrrole geochemistry, 547
 pigment content, gas-chromatography analyses, 547
 mass-spectrometric analyses, 547
 pigments, distribution with depth of burial, 547
 Thermal gradient, 87
 Canary Island volcanism effects, 70
 offshore wells, 70
 properties, fluids, 761
 sediments, 761
 Thermodynamic parameters, computer modeling, Site 397, 765
 Thermohaline bottom currents, equatorial Pacific, 783
 circulation in the Atlantic Ocean, intensification of, 36
 Thin-layer chromatography, 541
 Time-temperature effects, organic maturity, 506
 Titanomagnetite, volcaniclastic sandstones, 710
 Tortonian/Messinian boundary, foraminiflers, 339
 Site 397, 386
 Tracers of arid climates, quartz grains, 641
 Trade Wind belt off Cape Blanc, present influence of upwelling, 665
 Trade Winds in the Wheel-Round latitudes, paleo-winds, 737
 Transformation of dispersed organic matter (DOM), 584
 to opal-CT quartz, siliceous organisms, 742
 Transgression, Aaiun Basin, Late Cretaceous, 783
 Albian to Turonian, 19
 Transgressions, Middle Cretaceous, 811
 Transport and depositional mechanisms, organic matter in deep ocean sediments, 503
 Transverse debris flows, 84
Trapezopentus, new nannofossil genus, 232
 sarmatus, new nannofossil species, 232
 Trends, *n*-alkanes, 585
 Tripoli diatomites, 451
 Formation, 341
Triquetrorhabdulus carinatus Zone (NN 1), 291
 rugosus Zone, 380
 Triterpenes, 526
 Triterpenoid hydrocarbons, 516, 521, 542
 Trubi Formation at Capa Rossello, 341, 386
 of Sicily, 398
Tubodiscus jurapelicus Zone, 222
 Turbid layer transport, 564
 Turbidite sandstone beds, Site 397, middle to early Miocene, 806
 Turbidites, 36, 606, 610, 612, 616, 783, 797, 801
 in Canary Island Channel, 84
 middle to lower Miocene, 86
 Turbidity currents, 608, 611, 704
 Canary Island Channel, 715
 sediment transport by, 509
 Tyrrhenian Basin, Site 132, 457
 Sea, 382
 Unconformities, 491
 Unstable salinity regime, 585
 Uplift and shield stage, Canary Islands, early to middle Miocene of Canary Island Ridge, 790
 of Cape Verde Rise, 790
 Upper Cretaceous pelagic nannofossil marls, Site 369, 809
 and Paleogene sediments, Site 369, 221
 Miocene, sedimentation rate, 764
 Site 369, calcium carbonate dissolution 665
 Site 397, calcium carbonate dissolution, 665
 carbonate values, 672
 Neogene sedimentation rates, 751
 Quaternary influence, upwelling, Site 397, 665

- Upwelling, biogenic productivity effects, 611
 Early Miocene, 751
 along African continental margin south of Canary Islands, 798
 fish debris enrichment, 664
 influence on sediments, 664
 oxygen-isotope record effects, 735
 phosphorite occurrences, 664
 Site 397, present influence, 665
 upper Quaternary influence, 665
 Trade Wind belt off Cape Blanc, present influence, 665
 conditions, effect on biogenic productivity, 614
 maxima, opal-indicated, 735
 Variability in determinations of organic carbon, 506
 Variation downhole of natural remanent magnetization (NRM), Site 397, 475
 Viscosity, physical properties of fluids, 761
 Vitrinite reflectance, 505, 512, 805
 organic maturity, 518, 522
 values, 508
 type, organic matter, 506, 511, 514
 Vitrinite-huminite, reflectivity, 511
 Volcanic and sedimentary detritus from Fuerteventura Island, 33
 apron at southeastern foot, age of sediments, Canary Islands, 794
 ash layers, 781
 components source, Site 397, Fuerteventura Island 714
 Gran Canaria Island, 714
 debris, Canary Island, 613
 clastic origin of, Site 397, 714
 eruption on Tenerife Island, 85
 glass, 746
 material, eolian transport of, Site 369, 666
 Site 397, coarse fraction analyses, 658
 submarine transport of, Site 397, 666
 sandstones, carbonate cement, 713
 of the Emperor Seamounts, 748
 uplift of Canary Archipelago, Neogene, 713
 Volcaniclastic debris flows, 704
 flows from, Canary Islands, 616
 rock fragments, tachylite, 710
 sandstone, petrography of, 707
 sandstones, 32, 603, 610, 746, 775, 780
 igneous components, clinopyroxene, 710
 feldspar, 720
 olivine, 710
 titanomagnetite, 710
 igneous rock fragments, alkalic microgabbros, 710
 crystalline basalt, 710
 middle Miocene, petrochemistry of, 699
 petrography of, 699
 non-volcanic components, Crystalline rock fragments, 710
 quartz, 712
 sedimentary rock fragments, 710
 sedimentary structures, 28
 Site 397, 703
 zeolite cement, 713
 sediments, Site 397, alteration of, 746
 Volcanism, Canary Islands, 701
 coarse fraction analysis, 666
 middle Miocene to Quaternary, Sites 369 and 397, 796
 Volume of Mediterranean evaporite, 453
 Volume susceptibility, paleomagnetics, 69
 Vøring Plateau, organic-rich Miocene sediments from, 509
Watznaueria fasciata, new nannofossil species, 232
 Wealden facies, 19
 Early Cretaceous, 568
 sequence at the Cape Bojador margin, 810
 Wealden-like terrigenous sedimentation, 283
 Wealden-type delta foreset beds, Early Cretaceous, 807
 deltaic systems, 568, 810
 prograding delta system, 80
 Tan-Tan delta, 497
 West Saharan continental margin between Cape Bojador and Cape Blanc, Seismic reflection profiles of, 20
 margin, 21, 77
 Western African, continental rise, 727
 craton, 596
 Western Boundary countercurrents, 788
 X-ray diffraction studies, Lower Cretaceous silty clays, 579
 diffraction studies of sediments, Miocene/Pliocene boundary, 553
 results, clay minerals, 32
 studies, sediment diagenesis, 742
 of, Site 397 clay minerals, 595
 Zanclean/Piacenzian boundary, Site 397, 387
 Zeolite cement, volcaniclastic sandstones, 713
 Zeolites, 745, 746, 751
 cementation by, 747
 erionite, 747
 formation of, 87
 mordenite, 747
 Sites 397 and 398, 741